

Datasheet: 0200-0615 BATCH NUMBER 162221

| Description: | MOUSE ANTI HUMAN C3a |
|-------------------------|----------------------------|
| Specificity: | C3a |
| Other names: | COMPLEMENT COMPONENT 3a |
| Format: | Purified |
| | |
| Product Type: | Monoclonal Antibody |
| Product Type: Clone: | Monoclonal Antibody H13 |
| | - |

Product Details

| Applications | This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> . | | | | | |
|-----------------------------|--|------------------------|--------------|------------------------|---------------------|--|
| | | Yes | No | Not Determined | Suggested Dilution | |
| | Immunohistology - Frozen | • | | | 1/10 | |
| | Immunohistology - Paraffin | | | • | | |
| | ELISA | - | | | | |
| | Western Blotting | - | | | | |
| | Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given a a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls. | | | | | |
| Target Species | Human | | | | | |
| Species Cross Reactivity | Reacts with: Guinea Pig, N.B. Antibody reactivity a reactivity is derived from personal communications further information. | and worki testing w | ithin our la | boratories, peer-revie | wed publications or | |
| Product Form | Purified IgG - lyophilized | | | | | |
| Reconstitution | Reconstitute with 1.0ml of Care should be taken du bottom of the vial. Bio-Ra | ring recor | nstitution a | | | |

| Preparation | Purified IgG prepared by affinity chromatography on Protein A | | |
|-----------------------------------|---|--|--|
| Buffer Solution | Phosphate buffered saline | | |
| Preservative Stabilisers | 0.09% Sodium Azide, 0.5% Bovine Serum Albumin. | | |
| Approx. Protein Concentrations | IgG concentration 50 ug/ml after reconstitution. | | |
| Immunogen | Human complement component C3. | | |
| External Database Links | UniProt: <u>P01024</u> <u>Related reagents</u> Entrez Gene: | | |
| | 718 C3 Related reagents | | |
| Synonyms | CPAMD1 | | |
| RRID | AB_2195325 | | |
| Fusion Partners | Spleen cells from immunized Balb/c mice were fused with cells of the non-secreting P3-X63-Ag8.653 myeloma cell line. | | |
| | | | |
| Specificity | Mouse anti Human C3a antibody, clone H13 recognizes human C3a anaphylatoxin, a cleavage product of human complement component C3. C3a increases vascular permeability and causes histamine release from mast cells and basophilic leukocytes. | | |
| Specificity | cleavage product of human complement component C3. C3a increases vascular | | |
| Specificity | cleavage product of human complement component C3. C3a increases vascular permeability and causes histamine release from mast cells and basophilic leukocytes. Mouse anti Human C3a antibody, clone H13 recognizes a C3a-determinant also present on native C3. Mouse anti Human C3a antibody, clone H13 allows discrimination of ubiquitous or non-specifically bound native C3 or C3a from biologically active C3b fragment, neither C3b or C5 components are recognized by Mouse anti Human C3a | | |
| Specificity | cleavage product of human complement component C3. C3a increases vascular permeability and causes histamine release from mast cells and basophilic leukocytes. Mouse anti Human C3a antibody, clone H13 recognizes a C3a-determinant also present on native C3. Mouse anti Human C3a antibody, clone H13 allows discrimination of ubiquitous or non-specifically bound native C3 or C3a from biologically active C3b fragment, neither C3b or C5 components are recognized by Mouse anti Human C3a antibody, clone H13 (Burger <i>et al.</i> 1987). Mouse anti Human C3a antibody, clone H13 is reported to block C3a functional activity | | |

 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

 Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

 Guarantee
 Guaranteed until date of expiry. Please see product label.

 Health And Safety
 Material Safety Datasheet documentation #20486 available at: https://www.bio-rad-antibodies.com/SDS/0200-0615

 Regulatory
 For research purposes only

Related Products

Recommended Secondary Antibodies

| Rabbit Anti Mouse IgG (STAR12) | RPE |
|-------------------------------------|---|
| Goat Anti Mouse IgG IgA IgM (STAR87 |) <u>HRP</u> |
| Goat Anti Mouse IgG (STAR76) | RPE |
| Goat Anti Mouse IgG (STAR70) | <u>FITC</u> |
| Rabbit Anti Mouse IgG (STAR13) | HRP |
| Goat Anti Mouse IgG (Fc) (STAR120) | FITC, HRP |
| Rabbit Anti Mouse IgG (STAR9) | <u>FITC</u> |
| Goat Anti Mouse IgG (STAR77) | HRP |
| Goat Anti Mouse IgG (H/L) (STAR117) | <u>Alk. Phos.</u> , <u>DyLight®488</u> , <u>DyLight®550</u> , |
| | DyLight®650, DyLight®680, DyLight®800, |
| | FITC, HRP |

| North & South | Tel: +1 800 265 7376 | Worldwide | Tel: +44 (0)1865 852 700 | Europe | Tel: +49 (0) 89 8090 95 21 |
|---------------|---------------------------------|-----------|------------------------------|----------|--------------------------------------|
| America | Fax: +1 919 878 3751 | | Fax: +44 (0)1865 852 739 | | Fax: +49 (0) 89 8090 95 50 |
| | Email: antibody_sales_us@bio-ra | ad.com | Email: antibody_sales_uk@bio | -rad.com | Email: antibody_sales_de@bio-rad.com |

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M391273:211008'

Printed on 04 Mar 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint