

## Datasheet: 0300-0004

<b>Description:</b>	RABBIT ANTI RAT MDM2
<b>Specificity:</b>	MDM2
<b>Format:</b>	Purified
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	1 ml

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			■	
Immunohistology - Frozen		■		
Immunohistology - Paraffin		■		
Immunohistology - Resin		■		
ELISA	■			
Western Blotting			■	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

<b>Target Species</b>	Rat
<b>Species Cross Reactivity</b>	Reacts with: Horse Based on sequence similarity, is expected to react with: Mammals <b>N.B.</b> Antibody reactivity and working conditions may vary between species.
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on Protein G
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Approx. Protein Concentrations</b>	IgG concentration 5.0 mg/ml
<b>Immunogen</b>	Synthetic peptide corresponding to the amino acids 154-167 (C)SRPSTSSRRRAISE) of rat MDM2 conjugated to Keyhole Limpet Haemocyanin.

<b>Specificity</b>	<b>Rabbit anti Rat MDM2 antibody</b> recognizes rat MDM2, a nuclear phosphoprotein that controls intracellular levels of the tumour suppressor protein p53 by ubiquitination and subsequent proteasomal degradation. MDM2 also interacts with the retinoblastoma (RB) protein and E2F-1, to promote G1-S phase transition in the cell cycle.
<b>Storage</b>	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
<b>Shelf Life</b>	18 months from date of despatch.
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: 10040: <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</a>
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...)	<a href="#">FITC</a>
Sheep Anti Rabbit IgG (STAR35...)	<a href="#">RPE</a>
Goat Anti Rabbit IgG (H/L) (STAR124...)	<a href="#">HRP</a>
Goat Anti Rabbit IgG (Fc) (STAR121...)	<a href="#">Biotin</a> , <a href="#">FITC</a> , <a href="#">HRP</a>
Sheep Anti Rabbit IgG (2AB02...)	<a href="#">Biotin</a>
Sheep Anti Rabbit IgG (STAR36...)	<a href="#">DyLight@488</a> , <a href="#">DyLight@549</a> , <a href="#">DyLight@649</a> , <a href="#">DyLight@680</a> , <a href="#">DyLight@800</a>

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

'M308698:170807'

Printed on 25 May 2018