

Datasheet: 0300-0230 **BATCH NUMBER 160424**

Description:	RABBIT ANTI HUMAN C4d
Specificity:	C4d
Other names:	COMPLEMENT COMPONENT 4d
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.25 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	•			1/20 - 1/80
Immunohistology - Paraffin (1)	•			1/20 - 1/80
Immunofluorescence				1/20 - 1/80

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. TRIS/EDTA buffer pH 8.5 is recommended for this purpose.

Target Species	Human
Product Form	Purified Ig - liquid
Preparation	Purified Ig prepared by affinity chromatography on Protein G
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.1% Proclin™300
Immunogen	Synthetic peptide corresponding to amino acids 1241-1256 of C4 conjugated to Keyho limpet Haemocyanin.

External Database Links	UniProt: P0C0L4 Related reagents
	P0C0L5 Related reagents
	Entrez Gene:
	720 C4A Related reagents
	721 C4B Related reagents
Synonyms	CO4, CPAMD2, CPAMD3
RRID	AB_620313
Specificity	Rabbit anti Human C4d antibody is specific for human complement split product C4d.
References	 Regele, H. <i>et al.</i> (2001) Endothelial C4d deposition is associated with inferior kidney allograft outcome independently of cellular rejection. Nephrol Dial Transplant. 16 (10): 2058-66. Santos, A. <i>et al.</i> (2012) C4d detection in renal allograft biopsies: immunohistochemistry vs. immunofluorescence Port J Nephrol Hypert 26 (4): 272-7. Sethi, S. <i>et al.</i> (2015) C4d as a Diagnostic Tool in Proliferative GN. J Am Soc Nephrol. 26 (11): 2852-9. Viana, H. <i>et al.</i> (2009) C4d presence in kidney allograft biopsy: sensitivity and specificity of immunoperoxidase vs. immunofluorescence. N: 10th Banff Conference on Allograft Pathology, 2009, 9 a 14 Ago. Banff, Alberta, Canada Sreedharanunni, S. <i>et al.</i> (2014) An analysis of transplant glomerulopathy and thrombotic microangiopathy in kidney transplant biopsies. Transpl Int. 27 (8): 784-92. Hanaoka, K. <i>et al.</i> (2019) Benefits of a loading dose of tacrolimus on graft survival of kidney transplants in nonhuman primates Transplant Immunology. 52: 32-9. Mahakur, S. <i>et al.</i> (2018) Allo-specific immune response profiles indicative of acute rejection in kidney allografts using an <i>in vitro</i> lymphocyte culture-based model. Clin Exp Nephrol. 22 (2): 465-73.
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	Guaranteed until date of expiry. Please see product label.
Health And Safety Information	Material Safety Datasheet documentation #10199 available at: https://www.bio-rad-antibodies.com/SDS/0300-0230 10199
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

Goat Anti Rabbit IgG (H/L) (STAR124...) HRP

Sheep Anti Rabbit IgG (STAR35...) RPE

Recommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M382520:210513'

Printed on 12 Sep 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint