

Datasheet: 0300-0230 **BATCH NUMBER 166174** 

Description:	RABBIT ANTI HUMAN C4d
Specificity:	C4d
Other names:	COMPLEMENT COMPONENT 4d
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.25 ml

## **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	•			1/20 - 1/80
Immunohistology - Paraffin (1)	•			1/20 - 1/80
Immunofluorescence				1/20 - 1/80

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. TRIS/EDTA buffer pH 8.5 is recommended for this purpose.

Target Species	Human
Product Form	Purified Ig - liquid
Preparation	Purified Ig prepared by affinity chromatography on Protein G.
Buffer Solution	Phosphate buffered saline.
Preservative Stabilisers	0.1% Proclin 950
Immunogen	Synthetic peptide corresponding to amino acids 1241-1256 of C4 conjugated to Ke limpet Haemocyanin.

Links  UniProt: POC0L4 Related reagents POC0L5 Related reagents  Entrez Gene: T20 C4A Related reagents T21 C4B Related reagents T21 C4B Related reagents T21 C4B Related reagents T21 C4B Related reagents  Synonyms  CO4, CPAMD2, CPAMD3  RRID AB_620313  Specificity Rabbit anti Human C4d antibody is specific for human complement split product C4d.  References  1. Regele, H. et al. (2001) Endothelial C4d deposition is associated with inferior kidney allograft outcome independently of cellular rejection. Nephrol Dial Transplant. 16 (10): 2058-66. 2. Viana, H. et al. (2009) C4d presence in kidney allograft biopsy: sensitivity and specificity of immunoperoxidase vs. immunofluorescence. N: 10th Banff Conference on Allograft Pathology. 2009. 9 a 14 Aqo. Banff. Alberta, Canada 3. Santos, A. et al. (2012) C4d detection in renal allograft biopsies: immunohistochemistry vs. immunofluorescence Port J. Nephrol Hypert 26 (4): 272-7. 4. Sreedharanunni, S. et al. (2014) An analysis of transplant glomerulopathy and thrombotic microangiopathy in kidney transplant biopsies. Transpl. Int. 27 (8): 784-92. 5. Sethi, S. et al. (2015) C4d as a Diagnostic Tool in Proliferative GN. J.Am Soc Nephrol. 26 (11): 2852-9. 6. Mahakur, S. et al. (2018) Allo-specific immune response profiles indicative of acute rejection in kidney allografts using an <i>in vitro</i> lymphocyte culture-based model. Clin Exp. Nephrol. 22 (2): 465-73. 7. Hanaoka, K. et al. (2019) Benefits of a loading dose of tacrolimus on graft survival of kidney transplants in nonhuman primates Transplant Immunology. 52: 32-9.  Storage  This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.					
Entrez Gene:  720 C4A Related reagents 721 C4B Related reagents 721 C4B Related reagents 721 C4B Related reagents 721 C4B Related reagents 722 C4B Related reagents 723 Related reagents 724 C4B Related reagents 725 Related reagents 726 Related reagents 727 Repelle, H. et al. (2001) Endothelial C4d deposition is associated with inferior kidney allograft outcome independently of cellular rejection. Nephrol Dial Transplant. 16 (10): 2058-86. 72 Viana, H. et al. (2009) C4d presence in kidney allograft biopsy; sensitivity and specificity of immunoperoxidase vs. immunofluorescence. N. 10th Banff Conference on Allograft Pathology. 2009. 9 a 14 Ago. Banff, Alberta. Canada 73 Santos, A. et al. (2012) C4d detection in renal allograft biopsies: immunohistochemistry vs. immunofluorescence Port J Nephrol Hypert 26 (4): 272-7 74. Sreedharanunni, S. et al. (2014) An analysis of transplant glomerulopathy and thrombotic microangiopathy in kidney transplant biopsies. Transplant 26 (11): 2852-9. 75 Sethi, S. et al. (2015) C4d as a Diagnostic Tool in Proliferative GN. J Am Soc Nephrol. 26 (11): 2852-9. 86 Mahakur, S. et al. (2018) Allo-specific immune response profiles indicative of acute rejection in kidney allografts using an in vitro lymphocyte culture-based model. Clin Exp Nephrol. 22 (2): 465-73. 76 Hanaoka, K. et al. (2019) Benefits of a loading dose of tacrolimus on graft survival of kidney transplants in nonhuman primates Transplant Immunology. 52: 32-9.  Storage  This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.  Guarantee  Guaranteed until date of expiry, Please see product label.  Health And Safety Information antibodies.com/SDS/0300-0230	External Database Links	UniProt:			
Entrez Gene:  720 C4A Related reagents  721 C4B Related reagents  Synonyms CO4, CPAMD2, CPAMD3  RRID AB_620313  Specificity Rabbit anti Human C4d antibody is specific for human complement split product C4d.  References 1. Regele, H. et al. (2001) Endothelial C4d deposition is associated with inferior kidney allograft outcome independently of cellular rejection. Nephrol Dial Transplant. 16 (10): 2058-66.  2. Viana, H. et al. (2009) C4d presence in kidney allograft biopsy: sensitivity and specificity of immunoperoxidase vs. immunofluorescence. N: 10th Banff Conference on Allograft Pathology. 2009. 9 a 14 Ago. Banff. Alberta, Canada  3. Santos, A. et al. (2012) C4d detection in renal allograft biopsies: immunohistochemistry vs. immunofluorescence Port. Nephrol Hypert 26 (4): 272-7.  4. Sreedharanunni, S. et al. (2014) An analysis of transplant glomerulopathy and thrombotic microangiopathy in kidney transplant biopsies. Transpl Int. 27 (8): 784-92.  5. Sethi, S. et al. (2015) C4d as a Diagnostic Tool in Proliferative GN. J Am Soc Nephrol. 26 (111): 2852-9.  6. Mahakur, S. et al. (2018) Allo-specific immune response profiles indicative of acute rejection in kidney allografts using an in vitro lymphocyte culture-based model. Clin Exp. Nephrol. 22 (2): 465-73.  7. Hanaoka,K. et al. (2019) Benefits of a loading dose of tacrolimus on graft survival of kidney transplants in nonhuman primates Transplant Immunology. 52: 32-9.  Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.  Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.  Guarantee Guaranteed until date of expiry. Please see product label.  Health And Safety Information Material Safety Datasheet documentation #20479 available at: https://www.bio-rad-antibodies.com/SDS/0300-0230		P0C0L4 Related reagents			
Synonyms  CO4, CPAMD2, CPAMD3  RRID  AB_620313  Specificity  Rabbit anti Human C4d antibody is specific for human complement split product C4d.  References  1. Regele, H. et al. (2001) Endothelial C4d deposition is associated with inferior kidney allograft outcome independently of cellular rejection. Nephrol Dial Transplant. 16 (10): 2058-66.  2. Viana, H. et al. (2009) C4d presence in kidney allograft biopsy: sensitivity and specificity of immunoperoxidase vs. immunofluorescence. N: 10th Banff Conference on Allograft Patholocy. 2009. 9 a 14 Ago. Banff. Alberta. Canada  3. Santos, A. et al. (2012) C4d detection in renal allograft biopsies: immunohistochemistry vs. immunofluorescence Port J. Nephrol Hypert 26 (4): 272-7.  4. Sreedharanunni, S. et al. (2014) An analysis of transplant glomerulopathy and thrombotic microangiopathy in kidney transplant biopsies. Transpl Int. 27 (8): 784-92.  5. Sethi, S. et al. (2015) C4d as a Diagnostic Tool in Proliferative GN. J.Am Soc Nephrol. 26 (11): 2852-9.  6. Mahakur, S. et al. (2018) Allo-specific immune response profiles indicative of acute rejection in kidney allografts using an in vitro lymphocyte culture-based model. Clin Exp. Nephrol. 22 (2): 465-73.  7. Hanaoka, K. et al. (2019) Benefits of a loading dose of tacrolimus on graft survival of kidney transplants in nonhuman primates Transplant Immunology. 52: 32-9.  Storage  This product is shipped at ambient temperature. It is recommended to aliquot and store at 20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.  Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.  Guarantee  Guaranteed until date of expiry. Please see product label.  Health And Safety Internation #20479 available at: https://www.bio-rad-antibodies.com/SDS/0300-0230		P0C0L5 Related reagents			
Synonyms  CO4, CPAMD2, CPAMD3  RRID  AB_620313  Specificity  Rabbit anti Human C4d antibody is specific for human complement split product C4d.  1. Regele, H. et al. (2001) Endothelial C4d deposition is associated with inferior kidney allograft outcome independently of cellular rejection. Nephrol Dial Transplant. 16 (10): 2058-66.  2. Viana, H. et al. (2009) C4d presence in kidney allograft biopsy: sensitivity and specificity of immunoperoxidase vs. immunofluorescence. N: 10th Banff Conference on Allograft Pathology. 2009. 9 a 14 Aqo. Banff, Alberta, Canada  3. Santos, A. et al. (2012) C4d detection in renal allograft biopsies: immunohistochemistry vs. immunofluorescence Port J Nephrol Hypert 26 (4): 272-7.  4. Sreedharanunni, S. et al. (2014) An analysis of transplant glomerulopathy and thrombotic microangiopathy in kidney transplant biopsies. Transpl Int. 27 (8): 784-92.  5. Sethi, S. et al. (2015) C4d as a Diagnostic Tool in Proliferative GN. J Am Soc Nephrol. 26 (11): 2852-9.  6. Mahakur, S. et al. (2018) Allo-specific immune response profiles indicative of acute rejection in kidney allografts using an in vitro lymphocyte culture-based model. Clin Exp Nephrol. 22 (2): 465-73.  7. Hanaoka, K. et al. (2019) Benefits of a loading dose of tacrolimus on graft survival of kidney transplants in nonhuman primates Transplant Immunology. 52: 32-9.  Storage  This product is shipped at ambient temperature. It is recommended to aliquot and store a -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.  Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.  Guarantee  Guaranteed until date of expiry. Please see product label.  Health And Safety Intromation  Material Safety Datasheet documentation #20479 available at: https://www.bio-rad-antibodies.com/SDS/0300-0230		Entrez Gene:			
Synonyms  CO4, CPAMD2, CPAMD3  RRID  AB_620313  Specificity  Rabbit anti Human C4d antibody is specific for human complement split product C4d.  1. Regele, H. et al. (2001) Endothelial C4d deposition is associated with inferior kidney allograft outcome independently of cellular rejection. Nephrol Dial Transplant. 16 (10): 2058-66.  2. Viana, H. et al. (2009) C4d presence in kidney allograft biopsy: sensitivity and specificity of immunoperoxidase vs. immunofluorescence. N: 10th Banff Conference on Allograft Pathology. 2009. 9 a 14 Aqo. Banff, Alberta, Canada  3. Santos, A. et al. (2012) C4d detection in renal allograft biopsies: immunohistochemistry vs. immunofluorescence Port J Nephrol Hypert 26 (4): 272-7.  4. Sreedharanunni, S. et al. (2014) An analysis of transplant glomerulopathy and thrombotic microangiopathy in kidney transplant biopsies. Transpl Int. 27 (8): 784-92.  5. Sethi, S. et al. (2015) C4d as a Diagnostic Tool in Proliferative GN. J Am Soc Nephrol. 26 (11): 2852-9.  6. Mahakur, S. et al. (2018) Allo-specific immune response profiles indicative of acute rejection in kidney allografts using an in vitro lymphocyte culture-based model. Clin Exp Nephrol. 22 (2): 465-73.  7. Hanaoka, K. et al. (2019) Benefits of a loading dose of tacrolimus on graft survival of kidney transplants in nonhuman primates Transplant Immunology. 52: 32-9.  Storage  This product is shipped at ambient temperature. It is recommended to aliquot and store a -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.  Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.  Guarantee  Guaranteed until date of expiry. Please see product label.  Health And Safety Intromation  Material Safety Datasheet documentation #20479 available at: https://www.bio-rad-antibodies.com/SDS/0300-0230					
RRID AB_620313  Specificity Rabbit anti Human C4d antibody is specific for human complement split product C4d.  References 1. Regele, H. et al. (2001) Endothelial C4d deposition is associated with inferior kidney allograft outcome independently of cellular rejection. Nephrol Dial Transplant. 16 (10): 2058-66. 2. Viana, H. et al. (2009) C4d presence in kidney allograft biopsy: sensitivity and specificity of immunoperoxidase vs. immunofluorescence. N: 10th Banff Conference on Allograft Pathology. 2009. 9 a 14 Ago. Banff. Alberta. Canada 3. Santos, A. et al. (2012) C4d detection in renal allograft biopsies: immunohistochemistry vs. immunofluorescence Port J Nephrol Hypert 26 (4): 272-7. 4. Sreedharanunni, S. et al. (2014) An analysis of transplant glomerulopathy and thrombotic microangiopathy in kidney transplant biopsies. Transpl Int. 27 (8): 784-92. 5. Sethi, S. et al. (2015) C4d as a Diagnostic Tool in Proliferative GN. JAm Soc Nephrol. 26 (11): 2852-9. 6. Mahakur, S. et al. (2018) Allo-specific immune response profiles indicative of acute rejection in kidney allografts using an in vitro lymphocyte culture-based model. Clin Exp Nephrol. 22 (2): 465-73. 7. Hanaoka, K. et al. (2019) Benefits of a loading dose of tacrolimus on graft survival of kidney transplants in nonhuman primates Transplant Immunology. 52: 32-9.  Storage This product is shipped at ambient temperature. It is recommended to aliquot and store a -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.  Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.  Guarantee Guaranteed until date of expiry. Please see product label.  Material Safety Datasheet documentation #20479 available at: https://www.bio-rad-antibodies.com/SDS/0300-0230 20479					
RRID  AB_620313  Specificity  Rabbit anti Human C4d antibody is specific for human complement split product C4d.  References  1. Regele, H. et al. (2001) Endothelial C4d deposition is associated with inferior kidney allograft outcome independently of cellular rejection. Nephrol Dial Transplant. 16 (10): 2058-66.  2. Viana, H. et al. (2009) C4d presence in kidney allograft biopsy: sensitivity and specificity of immunoperoxidase vs. immunofluorescence. N: 10th Banff Conference on Allograft Pathology. 2009. 9 a 14 Ago. Banff. Alberta. Canada  3. Santos, A. et al. (2012) C4d detection in renal allograft biopsies: immunohistochemistry vs. immunofluorescence Port J Nephrol Hypert 26 (4): 272-7.  4. Sreedharanunni, S. et al. (2014) An analysis of transplant glomerulopathy and thrombotic microangiopathy in kidney transplant biopsies. Transpl Int. 27 (8): 784-92.  5. Sethi, S. et al. (2015) C4d as a Diagnostic Tool in Proliferative GN. J Am Soc Nephrol. 26 (11): 2852-9.  6. Mahakur, S. et al. (2018) Allo-specific immune response profiles indicative of acute rejection in kidney allografts using an in vitro lymphocyte culture-based model. Clin Exp Nephrol. 22 (2): 465-73.  7. Hanaoka, K. et al. (2019) Benefits of a loading dose of tacrolimus on graft survival of kidney transplants in nonhuman primates Transplant Immunology. 52: 32-9.  Storage  This product is shipped at ambient temperature. It is recommended to aliquot and store al -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.  Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.  Guarantee  Guaranteed until date of expiry. Please see product label.  Material Safety Datasheet documentation #20479 available at: https://www.bio-rad-antibodies.com/SDS/0300-0230 20479					
References  1. Regele, H. et al. (2001) Endothelial C4d deposition is associated with inferior kidney allograft outcome independently of cellular rejection. Nephrol Dial Transplant. 16 (10): 2058-66.  2. Viana, H. et al. (2009) C4d presence in kidney allograft biopsy: sensitivity and specificity of immunoperoxidase vs. immunofluorescence. N: 10th Banff Conference on Allograft Pathology. 2009. 9 a 14 Ago. Banff, Alberta. Canada 3. Santos, A. et al. (2012) C4d detection in renal allograft biopsies: immunohistochemistry vs. immunofluorescence Port J Nephrol Hypert 26 (4): 272-7. 4. Sreedharanunni, S. et al. (2014) An analysis of transplant glomerulopathy and thrombotic microangiopathy in kidney transplant biopsies. Transpl Int. 27 (8): 784-92. 5. Sethi, S. et al. (2015) C4d as a Diagnostic Tool in Proliferative GN. J Am Soc Nephrol. 26 (11): 2852-9. 6. Mahakur, S. et al. (2018) Allo-specific immune response profiles indicative of acute rejection in kidney allografts using an in vitro lymphocyte culture-based model. Clin Exp Nephrol. 22 (2): 465-73. 7. Hanaoka, K. et al. (2019) Benefits of a loading dose of tacrolimus on graft survival of kidney transplants in nonhuman primates Transplant Immunology. 52: 32-9.  Storage  This product is shipped at ambient temperature. It is recommended to aliquot and store a -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.  Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.  Guarantee  Guaranteed until date of expiry. Please see product label.  Material Safety Datasheet documentation #20479 available at: https://www.bio-rad-antibodies.com/SDS/0300-0230 20479	Synonyms	CO4, CPAMD2, CPAMD3			
1. Regele, H. et al. (2001) Endothelial C4d deposition is associated with inferior kidney allograft outcome independently of cellular rejection. Nephrol Dial Transplant. 16 (10): 2058-66.  2. Viana, H. et al. (2009) C4d presence in kidney allograft biopsy: sensitivity and specificity of immunoperoxidase vs. immunofluorescence. N: 10th Banff Conference on Allograft Pathology. 2009. 9 a 14 Ago. Banff, Alberta. Canada  3. Santos, A. et al. (2012) C4d detection in renal allograft biopsies: immunohistochemistry vs. immunofluorescence Port J Nephrol Hypert 26 (4): 272-7.  4. Sreedharanunni, S. et al. (2014) An analysis of transplant glomerulopathy and thrombotic microangiopathy in kidney transplant biopsies. Transpl Int. 27 (8): 784-92.  5. Sethi, S. et al. (2015) C4d as a Diagnostic Tool in Proliferative GN. J Am Soc Nephrol. 26 (11): 2852-9.  6. Mahakur, S. et al. (2018) Allo-specific immune response profiles indicative of acute rejection in kidney allografts using an in vitro lymphocyte culture-based model. Clin Exp Nephrol. 22 (2): 465-73.  7. Hanaoka, K. et al. (2019) Benefits of a loading dose of tacrolimus on graft survival of kidney transplants in nonhuman primates Transplant Immunology. 52: 32-9.  Storage  This product is shipped at ambient temperature. It is recommended to aliquot and store a -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.  Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.  Guarantee  Guaranteed until date of expiry. Please see product label.  Material Safety Datasheet documentation #20479 available at: https://www.bio-rad-antibodies.com/SDS/0300-0230 20479	RRID	AB_620313			
allograft outcome independently of cellular rejection. Nephrol Dial Transplant. 16 (10): 2058-66.  2. Viana, H. et al. (2009) C4d presence in kidney allograft biopsy: sensitivity and specificity of immunoperoxidase vs. immunofluorescence. N: 10th Banff Conference on Allograft Pathology. 2009. 9 a 14 Ago. Banff, Alberta. Canada  3. Santos, A. et al. (2012) C4d detection in renal allograft biopsies: immunohistochemistry vs. immunofluorescence Port J Nephrol Hypert 26 (4): 272-7.  4. Sreedharanunni, S. et al. (2014) An analysis of transplant glomerulopathy and thrombotic microangiopathy in kidney transplant biopsies. Transpl Int. 27 (8): 784-92.  5. Sethi, S. et al. (2015) C4d as a Diagnostic Tool in Proliferative GN. JAm Soc Nephrol. 26 (11): 2852-9.  6. Mahakur, S. et al. (2018) Allo-specific immune response profiles indicative of acute rejection in kidney allografts using an in vitro lymphocyte culture-based model. Clin Exp Nephrol. 22 (2): 465-73.  7. Hanaoka, K. et al. (2019) Benefits of a loading dose of tacrolimus on graft survival of kidney transplants in nonhuman primates Transplant Immunology. 52: 32-9.  Storage  This product is shipped at ambient temperature. It is recommended to aliquot and store a -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.  Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.  Guarantee  Guaranteed until date of expiry. Please see product label.  Material Safety Datasheet documentation #20479 available at: https://www.bio-rad-antibodies.com/SDS/0300-0230 20479	Specificity	Rabbit anti Human C4d antibody is specific for human complement split product C4d.			
-20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.  Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.  Guarantee  Guaranteed until date of expiry. Please see product label.  Health And Safety Information  Material Safety Datasheet documentation #20479 available at:  https://www.bio-rad-antibodies.com/SDS/0300-0230 20479	References	allograft outcome independently of cellular rejection. Nephrol Dial Transplant. 16 (10): 2058-66.  2. Viana, H. et al. (2009) C4d presence in kidney allograft biopsy: sensitivity and specificity of immunoperoxidase vs. immunofluorescence. N: 10th Banff Conference on Allograft Pathology, 2009, 9 a 14 Ago. Banff, Alberta, Canada  3. Santos, A. et al. (2012) C4d detection in renal allograft biopsies: immunohistochemistry vs. immunofluorescence Port J Nephrol Hypert 26 (4): 272-7.  4. Sreedharanunni, S. et al. (2014) An analysis of transplant glomerulopathy and thrombotic microangiopathy in kidney transplant biopsies. Transpl Int. 27 (8): 784-92.  5. Sethi, S. et al. (2015) C4d as a Diagnostic Tool in Proliferative GN. J Am Soc Nephrol. 26 (11): 2852-9.  6. Mahakur, S. et al. (2018) Allo-specific immune response profiles indicative of acute rejection in kidney allografts using an in vitro lymphocyte culture-based model. Clin Exp Nephrol. 22 (2): 465-73.  7. Hanaoka,K. et al. (2019) Benefits of a loading dose of tacrolimus on graft survival of			
Health And Safety Information  Material Safety Datasheet documentation #20479 available at:  https://www.bio-rad-antibodies.com/SDS/0300-0230 20479	Storage	-20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.  Avoid repeated freezing and thawing as this may denature the antibody. Storage in			
Information <a href="https://www.bio-rad-antibodies.com/SDS/0300-0230">https://www.bio-rad-antibodies.com/SDS/0300-0230</a> 20479	Guarantee	Guaranteed until date of expiry. Please see product label.			
Regulatory For research purposes only.	Health And Safety Information	https://www.bio-rad-antibodies.com/SDS/0300-0230			
	Regulatory	For research purposes only.			

# **Related Products**

## **Recommended Secondary Antibodies**

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

Sheep Anti Rabbit IgG (STAR35...) RPE

Goat Anti Rabbit IgG (H/L) (STAR124...) HRP

## **Recommended Useful Reagents**

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M419877:230622'

#### Printed on 29 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint