

Datasheet: 0300-0230 **BATCH NUMBER 167275**

| Description: | RABBIT ANTI HUMAN C4d |
|----------------------|-------------------------|
| Specificity: | C4d |
| Other names: | COMPLEMENT COMPONENT 4d |
| Format: | Purified |
| Product Type: | Polyclonal Antibody |
| Isotype: | Polyclonal IgG |
| Quantity: | 0.25 ml |
| | |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|--------------------------------|-----|----|----------------|--------------------|
| Immunohistology - Frozen | • | | | 1/20 - 1/80 |
| Immunohistology - Paraffin (1) | • | | | 1/20 - 1/80 |
| Immunofluorescence | | | | 1/20 - 1/80 |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

(1)This product requires antigen retrieval using heat treatment prior to staining of paraffin sections. TRIS/EDTA buffer pH 8.5 is recommended for this purpose.

| Target Species | Human |
|-----------------------------|---|
| Product Form | Purified Ig - liquid |
| Preparation | Purified Ig prepared by affinity chromatography on Protein G. |
| Buffer Solution | Phosphate buffered saline. |
| Preservative Stabilisers | 0.1% ProClin 950 |
| Immunogen | Synthetic peptide corresponding to amino acids 1241-1256 of C4 conjugated to Keyhol limpet Haemocyanin. |

| External Database Links | UniProt: | | | | |
|-------------------------------|---|--|--|--|--|
| LIIIKS | P0C0L4 Related reagents | | | | |
| | P0C0L5 Related reagents | | | | |
| | | | | | |
| | Entrez Gene: | | | | |
| | 720 C4A Related reagents | | | | |
| | 721 C4B Related reagents | | | | |
| Synonyms | CO4, CPAMD2, CPAMD3 | | | | |
| RRID | AB_620313 | | | | |
| Specificity | Rabbit anti Human C4d antibody is specific for human complement split product C4d. | | | | |
| References | 1. Regele, H. <i>et al.</i> (2001) Endothelial C4d deposition is associated with inferior kidney | | | | |
| | allograft outcome independently of cellular rejection. Nephrol Dial Transplant. 16 (10): | | | | |
| | <u>2058-66.</u> | | | | |
| | 2. Viana, H. et al. (2009) C4d presence in kidney allograft biopsy: sensitivity and | | | | |
| | specificity of immunoperoxidase vs. immunofluorescence. N: 10th Banff Conference on | | | | |
| | Allograft Pathology, 2009, 9 a 14 Ago. Banff, Alberta, Canada | | | | |
| | 3. Santos, A. <i>et al.</i> (2012) C4d detection in renal allograft biopsies: immunohistochemistry | | | | |
| | vs. immunofluorescence Port J Nephrol Hypert 26 (4): 272-7. | | | | |
| | 4. Sreedharanunni, S. <i>et al.</i> (2014) An analysis of transplant glomerulopathy and thrombotic microangiopathy in kidney transplant biopsies. <u>Transpl Int. 27 (8): 784-92.</u> | | | | |
| | 5. Sethi, S. <i>et al.</i> (2015) C4d as a Diagnostic Tool in Proliferative GN. <u>J Am Soc Nephrol.</u> | | | | |
| | 26 (11): 2852-9. | | | | |
| | 6. Mahakur, S. <i>et al.</i> (2018) Allo-specific immune response profiles indicative of acute | | | | |
| | rejection in kidney allografts using an <i>in vitro</i> lymphocyte culture-based model. <u>Clin Exp</u> | | | | |
| | Nephrol. 22 (2): 465-73. | | | | |
| | 7. Hanaoka,K. <i>et al.</i> (2019) Benefits of a loading dose of tacrolimus on graft survival of | | | | |
| | kidney transplants in nonhuman primates <u>Transplant Immunology. 52: 32-9.</u> | | | | |
| Stavana | | | | | |
| Storage | This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for | | | | |
| | short term use (up to 4 weeks) and store the remaining aliquots at -20°C. | | | | |
| | short term use (up to 4 weeks) and store the remaining alliquots at -20 C. | | | | |
| | Avoid repeated freezing and thawing as this may denature the antibody. Storage in | | | | |
| | frost-free freezers is not recommended. | | | | |
| Guarantee | Guaranteed until date of expiry. Please see product label. | | | | |
| Acknowledgements | ProClin is a trademark of The Dow Chemical Company ("Dow") or an affiliated company of Dow. | | | | |
| Hoolth And C-f-tr | Matarial Cafety Datashart dansum anti-tion #20470 and Hall | | | | |
| Health And Safety Information | Material Safety Datasheet documentation #20479 available at: | | | | |
| | https://www.bio-rad-antibodies.com/SDS/0300-0230 20479 | | | | |
| | ZU413 | | | | |

Regulatory For research purposes only.

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) FITC

Goat Anti Rabbit IgG (H/L) (STAR124...) HRP

Sheep Anti Rabbit IgG (STAR35...) RPE

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

Recommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M423160:231006'

Printed on 29 Feb 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint