

## Datasheet: 101008

<b>Description:</b>	GOAT ANTI MOUSE IgG/A/M (H/L):Biotin
<b>Specificity:</b>	IgG IgA IgM (H/L)
<b>Format:</b>	Biotin
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			1/2000 - 1/10,000
Immunoprecipitation			▪	
Immunoblotting	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Product Form</b>	Ig Fraction conjugated to Biotin - liquid
<b>Antiserum Preparation</b>	Antisera to Mouse IgG, IgA and IgM were raised by repeated immunisation of goats with pooled mouse IgG, IgA and IgM paraproteins.
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	<0.1% Sodium Azide (NaN <sub>3</sub> )
<b>Approx. Protein Concentrations</b>	Ig concentration 0.5 mg/ml

## External Database

## Links

## UniProt:

<a href="#">P01864</a>	<a href="#">Related reagents</a>
<a href="#">P01872</a>	<a href="#">Related reagents</a>
<a href="#">P03987</a>	<a href="#">Related reagents</a>
<a href="#">P01873</a>	<a href="#">Related reagents</a>
<a href="#">P01868</a>	<a href="#">Related reagents</a>
<a href="#">P01865</a>	<a href="#">Related reagents</a>
<a href="#">P01869</a>	<a href="#">Related reagents</a>
<a href="#">P01878</a>	<a href="#">Related reagents</a>
<a href="#">P01867</a>	<a href="#">Related reagents</a>
<a href="#">P01863</a>	<a href="#">Related reagents</a>
<a href="#">P01844</a>	<a href="#">Related reagents</a>
<a href="#">P01843</a>	<a href="#">Related reagents</a>
<a href="#">P01845</a>	<a href="#">Related reagents</a>
<a href="#">P01834</a>	<a href="#">Related reagents</a>

## Entrez Gene:

<a href="#">380793</a>	Igh-1a	<a href="#">Related reagents</a>
<a href="#">16019</a>	Ighm	<a href="#">Related reagents</a>
<a href="#">16017</a>	Ighg1	<a href="#">Related reagents</a>
<a href="#">16019</a>	Ighm	<a href="#">Related reagents</a>
<a href="#">380793</a>	Igh-1a	<a href="#">Related reagents</a>
<a href="#">380795</a>	AI324046	<a href="#">Related reagents</a>
<a href="#">3514</a>	IGKC	<a href="#">Related reagents</a>
<a href="#">16016</a>	Ighg2b	<a href="#">Related reagents</a>
<a href="#">16017</a>	Ighg1	<a href="#">Related reagents</a>
<a href="#">16061</a>	Igh-VJ558	<a href="#">Related reagents</a>
<a href="#">110786</a>	Iglc2	<a href="#">Related reagents</a>
<a href="#">110787</a>	Iglc3	<a href="#">Related reagents</a>
<a href="#">380793</a>	Igh-1a	<a href="#">Related reagents</a>
<a href="#">433053</a>	LOC433053	<a href="#">Related reagents</a>

---

## Synonyms

Igh-4

---

## RRID

AB\_609675

---

## Specificity

**Goat anti Mouse IgG/A/M** reacts with the heavy and light chains of all major classes of mouse immunoglobulin.

The antibody has been adsorbed against human serum to minimise cross-reactivity with human immunoglobulins.

---

**References**

1. Ding, G. *et al.* (2010) Effect of cryopreservation on biological and immunological properties of stem cells from apical papilla. [J Cell Physiol. 22: 415-22.](#)
2. Shao, L. *et al.* (2015) Comparative *In Vitro* and *In Vivo* Studies of Porcine Rotavirus G9P[13] and Human Rotavirus Wa G1P[8]. [J Virol. 90 \(1\): 142-51.](#)
3. Iver, S. *et al.* (2019) Probing BAK and BAX Activation and Pore Assembly with Cytochrome c Release, Limited Proteolysis, and Oxidant-Induced Linkage. [Methods Mol Biol. 1877: 201-16.](#)

---

**Storage**

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee**

Guaranteed until date of expiry. Please see product label.

---

**Health And Safety Information**

Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

---

**Regulatory**

For research purposes only

---

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M387464:210629'

**Printed on 21 Mar 2022**