

Datasheet: 102009

BATCH NUMBER 164394

Description:	GOAT ANTI MOUSE IgM:RPE
Specificity:	IgM
Format:	RPE
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse		
Product Form	Purified Ig conjugated to R. Phycoerythrin (RPE) - liquid		
Max Ex/Em	Fluorophore RPE 488nm laser	Excitation Max (nm) 496	Emission Max (nm)

Antiserum Preparation Purified Ig prepared by affinity chromatography on IgM covalently linked to agarose

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃) Stabilizing agent (sucrose)
Approx. Protein Concentrations	lg concentration 0.5 mg/ml
Immunogen	Mouse IgM paraproteins

External Database Links	UniProt: P01872 Related reagents P01873 Related reagents		
	Entrez Gene: 16019 Ighm Related reagents 16019 Ighm Related reagents		
RRID	AB_616863		
Specificity	Goat anti mouse IgM antibody recognises the heavy chain of mouse IgM as demonstrated by ELISA and flow cytometry. Minimal cross reactivity is observed with human immunoglobulins.		
	Goat anti mouse IgM antibody has been cross absorbed against Mouse Ig G_1 , Ig G_{2a} , Ig G_{2b} , Ig G_3 and IgA, pooled human sera and purified human paraproteins.		
References	 Kerr, K. et al. (2010) Inflammatory cytokine responses in a pregnant mouse model of <i>Chlamydophila abortus</i> infection. Vet Microbiol. 144 (3-4): 392-8. Kamat, M.M. et al. (2016) Changes in Myeloid Lineage Cells in the Uterus and Peripheral Blood of Dairy Heifers During Early Pregnancy. Biol Reprod. 95 (3): 68. Monzo, H.J. et al. (2017) Insulin promotes cell migration by regulating PSA-NCAM. Exp Cell Res. 355 (1): 26-39. Desancé, M. et al. (2018) Chondrogenic Differentiation of Defined Equine Mesenchymal Stem Cells Derived from Umbilical Cord Blood for Use in Cartilage Repair Therapy. Int J Mol Sci. 19 (2)Feb 10 [Epub ahead of print]. Penadés, M. et al. (2018) Long-term implications of feed energy source in different genetic types of reproductive rabbit females. II. Immunologic status. Animal. 12 (9): 1877-85. Penadés, M. et al. (2019) Early deviations in performance, metabolic and immunological indicators affect stayability in rabbit females. Animal.: 1-10. 		
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.		
Guarantee	Guaranteed for 12 months from the date of despatch or until the date of expiry, whichever comes first. Please see label for expiry date.		

Material Safety Datasheet documentation #10045 available at:

https://www.bio-rad-antibodies.com/SDS/102009

Health And Safety

10045

For research purposes only

Information

Regulatory

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21
 To

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50
 find a

 Email: antibody_sales_us@bio-rad.com
 Email: antibody_sales_de@bio-rad.com
 Email: antibody_sales_de@bio-rad.com

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M402118:220718'

Printed on 25 Apr 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint