

Datasheet: 103001

**BATCH NUMBER 158773**

<b>Description:</b>	GOAT ANTI MOUSE IgG
<b>Specificity:</b>	IgG
<b>Format:</b>	Purified
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			1/2000 - 1/4000
Immunoprecipitation			▪	
Western Blotting			▪	
Immunoblotting	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Product Form</b>	Ig fraction - liquid
<b>Preparation</b>	Purified Ig prepared by affinity chromatography on pooled mouse IgG covalently linked to agarose
<b>Antiserum Preparation</b>	Antisera to mouse IgG were raised by repeated immunisations of goats with highly purified antigen.
<b>Buffer Solution</b>	Borate buffered saline
<b>Preservative</b>	None Present

## Stabilisers

---

Approx. Protein Concentrations	Ig concentration 1.0 mg/ml
--------------------------------	----------------------------

---

## External Database Links

### UniProt:

<a href="#">P01867</a>	<a href="#">Related reagents</a>
<a href="#">P01865</a>	<a href="#">Related reagents</a>
<a href="#">P01863</a>	<a href="#">Related reagents</a>
<a href="#">P01864</a>	<a href="#">Related reagents</a>
<a href="#">P01868</a>	<a href="#">Related reagents</a>
<a href="#">P01869</a>	<a href="#">Related reagents</a>
<a href="#">P03987</a>	<a href="#">Related reagents</a>

### Entrez Gene:

<a href="#">16016</a>	Ighg2b	<a href="#">Related reagents</a>
<a href="#">380793</a>	Igh-1a	<a href="#">Related reagents</a>
<a href="#">16017</a>	Ighg1	<a href="#">Related reagents</a>
<a href="#">16017</a>	Ighg1	<a href="#">Related reagents</a>
<a href="#">380793</a>	Igh-1a	<a href="#">Related reagents</a>
<a href="#">380793</a>	Igh-1a	<a href="#">Related reagents</a>
<a href="#">380795</a>	AI324046	<a href="#">Related reagents</a>

---

Synonyms	Igh-4
----------	-------

---

RRID	AB_609693
------	-----------

---

Specificity	<b>Goat anti Mouse IgG antibody</b> recognizes mouse IgG, recognising the heavy chain of mouse IgG1, IgG2a, IgG2b and IgG3 as demonstrated by ELISA.
-------------	--

Goat anti Mouse IgG antibody has been cross-adsorbed against mouse IgM, mouse IgA and human serum to reduce potential cross-reactivity.

## References

1. Joimel, U. *et al.* (2010) Stimulation of angiogenesis resulting from cooperation between macrophages and MDA-MB-231 breast cancer cells: proposed molecular mechanism and effect of tetrathiomolybdate. [BMC Cancer. 10: 375.](#)
2. Childs K *et al.* (2012) Paramyxovirus V proteins interact with the RNA Helicase LGP2 to inhibit RIG-I-dependent interferon induction. [J Virol. 86 \(7\): 3411-21.](#)
3. Moalli, F. *et al.* (2015) Intravital and whole-organ imaging reveals capture of melanoma-derived antigen by lymph node subcapsular macrophages leading to widespread deposition on follicular dendritic cells. [Front Immunol. 6: 114.](#)
4. Ramos-Sevillano, E. *et al.* (2016) PSGL-1 on Leukocytes is a Critical Component of the Host Immune Response against Invasive Pneumococcal Disease. [PLoS Pathog. 12 \(3\): e1005500.](#)
5. Abbate, F. *et al.* (2016) Acid-sensing ion channel immunoreactivities in the cephalic neuromasts of adult zebrafish. [Ann Anat. 207: 27-31.](#)

6. Shea, G.K. *et al.* (2020) Juxtacrine signalling via Notch and ErbB receptors in the switch to fate commitment of bone marrow-derived Schwann cells. [Eur J Neurosci. 52 \(5\): 3306-21.](#)

<b>Storage</b>	Store at -20°C only. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
<b>Guarantee</b>	Guaranteed until date of expiry. Please see product label.
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10123 available at: <a href="https://www.bio-rad-antibodies.com/SDS/10300110123">https://www.bio-rad-antibodies.com/SDS/10300110123</a>
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) [FITC](#), [HRP](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M350076:190307'

Printed on 14 Mar 2024