

Datasheet: 103001F BATCH NUMBER 166185

Description:	GOAT ANTI MOUSE IgG:FITC
Specificity:	lgG
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u>						
	rad-antibodies.com/protocols. Yes No Not Determined Suggested Dilution						
	Flow Cytometry			not Dotoniniou			
	Immunofluorescence	•					
	Where this product has	s not been te	sted for	use in a particular te	echnique this does not		
	necessarily exclude its use in such procedures. Suggested working dilutions are given as						
	a guide only. It is recor system using appropria	nmended tha	it the use	er titrates the produc			
Target Species	Mouse						
Product Form	Purified IgG conjugate	d to Fluoresc	ein Isoth	iocyanate Isomer 1	(FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation M	ax (nm)	Emission Max (nm)	_		
	FITC	490		525	_		
Antiserum Preparation	n Antisera to mouse IgG purified antigen. Purifie IgG covalently linked to	ed IgG was p	• •		of goats with highly graphy on pooled mouse		
Buffer Solution	Phosphate buffered sa	line					
Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃)					
Approx. Protein Concentrations	IgG concentration 1.0 r	ng/ml					

External Database						
Links	UniProt:					
	<u>P01867</u>	Related	reagents			
	<u>P01865</u>	<u>Related</u>	reagents			
	<u>P01863</u>	P01863 Related reagents				
	<u>P01864</u>	<u>Related</u>	reagents			
	P01868 Related reagents					
	<u>P01869</u>		l reagents			
	<u>P03987</u>	Related	l reagents			
	Entrez Ge	ne:				
	<u>16016</u>	lghg2b	Related reagents			
	<u>380793</u>	lgh-1a	Related reagents			
	<u>16017</u>	lghg1	Related reagents			
	<u>16017</u>	lghg1	Related reagents			
	<u>380793</u>	lgh-1a	Related reagents			
	<u>380793</u>	lgh-1a	Related reagents			
	<u>380795</u>	AI324046	Related reagents			
Synonyms	lgh-4					
Specificity	mouse IgG Goat anti M	1, IgG2a, Ig louse IgG al	antibody recognizes mouse IgG, recognising the heavy chain of G2b and IgG3 as demonstrated by ELISA. ntibody has been cross-adsorbed against mouse IgM, mouse IgA educe potential cross-reactivity.			
References	macrophag effect of tet 2. Childs K inhibit RIG- 3. Moalli, F derived ant deposition	es and MDA rathiomolybe et al. (2012 I-dependent c et al. (2015 tigen by lymp on follicular	 0) Stimulation of angiogenesis resulting from cooperation between A-MB-231 breast cancer cells: proposed molecular mechanism and date. <u>BMC Cancer. 10: 375.</u>) Paramyxovirus V proteins interact with the RNA Helicase LGP2 to t interferon induction. <u>J Virol. 86 (7): 3411-21.</u> 5) Intravital and whole-organ imaging reveals capture of melanoma-oh node subcapsular macrophages leading to widespread dendritic cells. <u>Front Immunol. 6: 114.</u> <i>et al.</i> (2016) PSGL-1 on Leukocytes is a Critical Component of the 			
	Host Immu <u>e1005500.</u> 5. Abbate, neuromasta 6. Shea, G switch to fa <u>3306-21.</u> 7. Siqueira	ne Respons F. <i>et al.</i> (201 s of adult zel .K. <i>et al.</i> (20 ate commitm , R.F. & Ferr	 e against Invasive Pneumococcal Disease. <u>PLoS Pathog. 12 (3)</u>: 16) Acid-sensing ion channel immunoreactivities in the cephalic brafish. <u>Ann Anat. 207: 27-31.</u> 20) Juxtacrine signalling via Notch and ErbB receptors in the ent of bone marrow-derived Schwann cells. <u>Eur J Neurosci. 52 (5)</u>: mandes, R.L. (2018) Cryopreservation of lymphocytes for in horses Pesquisa Veterinária Brasileira. 38 (11): 2019-22. 			
Storage	Host Immu <u>e1005500.</u> 5. Abbate, neuromasts 6. Shea, G switch to fa <u>3306-21.</u> 7. Siqueira immunolog	ne Respons F. <i>et al.</i> (201 s of adult zel .K. <i>et al.</i> (20 ate commitm , R.F. & Ferr ical studies i	 a) Acid-sensing ion channel immunoreactivities in the cephalic brafish. <u>Ann Anat. 207: 27-31.</u> 20) Juxtacrine signalling via Notch and ErbB receptors in the ent of bone marrow-derived Schwann cells. <u>Eur J Neurosci. 52 (5):</u> 			

-20°C on receipt. When thawed, aliquot the sample as need short term use (up to 4 weeks) and store the remaining aliqu Avoid repeated freezing and thawing as this may denature t frost-free freezers is not recommended. This product is pho					uots at -20°C. he antibody. Storage in	
Guarantee Guaranteed for 12 months from the date of despatch or until comes first. Please see label for expiry date.					the date of expiry, whicheve	
		sheet documentation #10 ntibodies.com/SDS/1030	at:			
Regulatory For research purposes		es only				
Iorth & South	Tel: +1 800 265 7 Fax: +1 919 878		Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody sales uk@bid	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody sales de@bio-rad.con	

Printed on 14 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint