

Datasheet: 103008

**BATCH NUMBER 166229**

<b>Description:</b>	GOAT ANTI MOUSE IgG:Biotin
<b>Specificity:</b>	IgG
<b>Format:</b>	Biotin
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
ELISA	▪			1/5,000 - 1/20,000
Immunoprecipitation			▪	
Western Blotting			▪	
Immunofluorescence	▪			
Immunoblotting	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

<b>Target Species</b>	Mouse
<b>Product Form</b>	Purified IgG conjugated to Biotin - liquid
<b>Preparation</b>	Purified IgG prepared by affinity chromatography on pooled mouse IgG covalently linked to agarose
<b>Antiserum Preparation</b>	Antisera to mouse IgG were raised by repeated immunisations of goats with highly purified antigen.
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	<0.1% Sodium Azide (NaN <sub>3</sub> )

---

**Approx. Protein Concentrations**Ig concentration 0.5 mg/ml

---

**External Database Links****UniProt:**

[P01867](#) [Related reagents](#)  
[P01865](#) [Related reagents](#)  
[P01863](#) [Related reagents](#)  
[P01864](#) [Related reagents](#)  
[P01868](#) [Related reagents](#)  
[P01869](#) [Related reagents](#)  
[P03987](#) [Related reagents](#)

**Entrez Gene:**

[16016](#) Ighg2b [Related reagents](#)  
[380793](#) Igh-1a [Related reagents](#)  
[16017](#) Ighg1 [Related reagents](#)  
[16017](#) Ighg1 [Related reagents](#)  
[380793](#) Igh-1a [Related reagents](#)  
[380793](#) Igh-1a [Related reagents](#)  
[380795](#) AI324046 [Related reagents](#)

---

**Synonyms**Igh-4

---

**RRID**AB\_2103446

---

**Specificity**

**Goat anti Mouse IgG antibody** recognizes mouse IgG, recognising the heavy chain of mouse IgG1, IgG2a, IgG2b and IgG3 as demonstrated by ELISA.

Goat anti Mouse IgG antibody has been cross-adsorbed against mouse IgM, mouse IgA and human serum to reduce potential cross-reactivity.

---

**References**

1. Joimel, U. *et al.* (2010) Stimulation of angiogenesis resulting from cooperation between macrophages and MDA-MB-231 breast cancer cells: proposed molecular mechanism and effect of tetrathiomolybdate. [BMC Cancer. 10: 375.](#)
2. Childs K *et al.* (2012) Paramyxovirus V proteins interact with the RNA Helicase LGP2 to inhibit RIG-I-dependent interferon induction. [J Virol. 86 \(7\): 3411-21.](#)
3. Moalli, F. *et al.* (2015) Intravital and whole-organ imaging reveals capture of melanoma-derived antigen by lymph node subcapsular macrophages leading to widespread deposition on follicular dendritic cells. [Front Immunol. 6: 114.](#)
4. Ramos-Sevillano, E. *et al.* (2016) PSGL-1 on Leukocytes is a Critical Component of the Host Immune Response against Invasive Pneumococcal Disease. [PLoS Pathog. 12 \(3\): e1005500.](#)
5. Abbate, F. *et al.* (2016) Acid-sensing ion channel immunoreactivities in the cephalic neuromasts of adult zebrafish. [Ann Anat. 207: 27-31.](#)
6. Shea, G.K. *et al.* (2020) Juxtacrine signalling via Notch and ErbB receptors in the switch to fate commitment of bone marrow-derived Schwann cells. [Eur J Neurosci. 52 \(5\):](#)

[3306-21.](#)

7. Siqueira, R.F. & Fernandes, R.L. (2018) Cryopreservation of lymphocytes for immunological studies in horses [Pesquisa Veterinária Brasileira. 38 \(11\): 2019-22.](#)

---

**Storage** This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee** Guaranteed for 12 months from the date of despatch or until the date of expiry, whichever comes first. Please see label for expiry date.

---

**Health And Safety Information** Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/103008>  
10040

---

**Regulatory** For research purposes only

---

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M420295:230706'

**Printed on 14 Mar 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)