

Datasheet: 1065002S BATCH NUMBER 158473

Description:	MOUSE ANTI HUMAN ZAP 70:FITC		
Specificity:	ZAP 70		
Format:	FITC		
Product Type:	Monoclonal Antibody		
Clone:	SBZAP		
lsotype:	lgG1		
Quantity:	25 µg		

Product Details

Applications	This product has been reported to work in the following applications. This information i				
	derived from testing within our laboratories, peer-reviewed publications or personal				
	communications from the originators. Please refer to references indicated for further				
	information. For general protocol recommendations, please visit <u>www.bio-</u>				
	rad-antibodies.com/protocols.				
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry (1)	-			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)Membrane permeabilisation is required for this application. Bio-Rad recommends the use of Leucoperm[™] (Product Code <u>BUF09</u>) for this purpose.

Target Species	Human			
Product Form	Purified IgG conjug	gated to Fluorescein Isoth	niocyanate Isomer 1	
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)	
	FITC	490	525	
Buffer Solution	Phosphate buffered saline			
Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃)			
Approx. Protein Concentrations	IgG concentration	0.1 mg/ml		

Immunogen	Synthetic peptide corresponding to amino acids 280-309 of human ZAP70 conjugated to keyhole limpet hemocyanin (KLH).
External Database Links	UniProt: P43403 Related reagents Entrez Gene: 7535 ZAP70 Related reagents
Synonyms	SRK
RRID	AB_616761
Specificity	Mouse anti Human ZAP 70 antibody, clone SBZAP recognizes human ZAP-70, a 70kDa tyrosine protein kinase which associates with the T-cell receptor (TCR) zeta chain and undergoes phosphorylation following TCR stimulation. ZAP-70 is primarily expressed in T lymphocytes and natural killer cells, where it plays a key role in T-cell receptor (TCR) signalling. Recent studies suggest that ZAP-70 is also expressed in a population of normal immature B-cells and in B-cells from a sub-set of patients with chronic lymphocytic leukaemia.
References	 Roifman, C.M. <i>et al.</i> (1989) Depletion of CD8+ cells in human thymic medulla results in selective immune deficiency. <u>J Exp Med. 170 (6): 2177-82.</u> Arpaia, E. <i>et al.</i> (1994) Defective T cell receptor signaling and CD8+ thymic selection in humans lacking zap-70 kinase. <u>Cell. 76 (5): 947-58.</u>
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.
Guarantee	Guaranteed until date of expiry. Please see product label.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/1065002S 10040
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:FITC (MCA928F)

Recommended Useful Reagents

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@t	pio-rad.com	Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M387160:210615'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint