

Datasheet: 1351102

Description:	READIDROP™ 7-AAD
Name:	READIDROP™ 7-AAD
Format:	Reagent
Product Type:	Accessory Reagent
Quantity:	3 x 3ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/250 - 1/500

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Product Information

7-aminoactinomycin (7-AAD) is a fluorescent dye that has a high affinity for GC-rich regions of double stranded DNA. 7-AAD is not membrane permeable and therefore requires the compromised membranes of dead cells to bind to DNA. Upon excitation with an appropriate light source, dead cells exhibit higher fluorescence intensity than live cells, thereby enabling clear discrimination of these populations by flow cytometry. Dead cells can then be excluded from further analysis and also sorted from cell populations of interest. The spectral properties (Excitation 546 nm/ Emission 647 nm) enable use in both single color and multi color flow experiments. The convenient dropper bottle format removes the manual weighing, pipetting and dilution steps required by other commonly available 7-AAD products.

Instructions For Use

Caution: Wear gloves, safety glasses/face protection, and protective clothing to avoid direct contact of REDI-Drop solutions with skin and eyes.

1. Resuspend 1×10^6 cells in 500 μ l flow cytometry buffer (1x phosphate buffered saline with 3% BSA). The cell concentration is now 2×10^6 /ml.
2. Add one to two drop of REDI-Drop 7-AAD to the cell suspension.
3. Vortex briefly.

4. Incubate for 1-10 min at room temperature.

Note: Adding one drop to a 500 µl cell suspension produces a final dye concentration of 1 µg/ml. This is typically sufficient to stain 1×10^6 cells. If one drop is not sufficient for your cell type, adding a second drop may improve staining efficacy. When working with a larger volume of cells, scale ReadiDrop solution amount accordingly.

5. Analyze cells by flow cytometry.

- Detect 7-AAD signal in FL3 on Bio-Rad's S3e Cell Sorter. Otherwise, select the most appropriate FL channel on your particular instrument.

References	1. Al Tanoury, Z. <i>et al.</i> (2020) Differentiation of the human PAX7-positive myogenic precursors/satellite cell lineage <i>in vitro</i> . Development. 147 (12): dev187344.
Storage	Store at room temperature
Guarantee	Guaranteed until date of expiry. Please see product label.
Health And Safety Information	Material Safety Datasheet documentation #1351102 available at: https://www.bio-rad-antibodies.com/SDS/1351102 1351102
Regulatory	For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M374490:201109'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)