

Datasheet: 1406-1460 BATCH NUMBER 165780

RABBIT ANTI HUMAN BMP-9
BMP-9
GROWTH DIFFERENTIATION FACTOR 2
Purified
Polyclonal Antibody
Polyclonal IgG
1 ml

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal							
	communications from the	refer to references indicated for further						
	information. For general protocol recommendations, please visit <u>www.bio-</u>							
	rad-antibodies.com/proto							
		Yes	No	Not Determined	Suggested Dilution			
	Immunohistology - Frozen							
	Immunohistology - Paraffin	-						
	ELISA	-			1/2000 - 1/20000			
	Western Blotting	-						
	Immunofluorescence	-						
	Where this product has n	ot been to	ested for	use in a particular tech	nique this does not			
	necessarily exclude its us	se in such	n procedu	res. Suggested workin	ng dilutions are given as			
	a guide only. It is recommended that the user titrates the product for use in the system using the appropriate negative/positive controls.							
Target Species	Human							
Product Form	Purified IgG - liquid							
Preparation	Purified IgG prepared by							
Buffer Solution	Phosphate buffered salin							
Preservative Stabilisers	0.09% Sodium Azide (Na							
Approx. Protein Concentrations	5.0mg/ml							

Immunogen	Synthetic peptide corresponding to the amino acids 87 - 98 (CKDDMGVPTLKYH) of the native molecule conjugated to keyhole limpet haemocyanin.
External Database Links	UniProt: <u>Q9UK05</u> <u>Related reagents</u> Entrez Gene: <u>2658</u> GDF2 <u>Related reagents</u>
Synonyms	BMP9
RRID	AB_2111086
Specificity	Rabbit anti Human BMP-9 antibody recognizes Human BMP9 (Bone Morphogenic Protein), a member of the TGF-beta superfamily.
	BMPs act through binding with a receptor complex consisting of type I and type II serine/threonine kinases, resulting ultimately in the activation of the Smad protein and mitogen-activated protein kinase (MAPK) signaling pathways. Several antagonist proteins, including, noggin, chordin, gremlin and follistatin, are responsible for modulating the signaling effects of BMPs, through the binding and blocking of receptor ligands, thereby preventing activation.
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/1406-1460 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (Fc) (STAR121...)Biotin, FITC, HRPSheep Anti Rabbit IgG (STAR35...)RPEGoat Anti Rabbit IgG (H/L) (STAR124...)HRPRecommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)

TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	d.com	Email: antibody sales uk@bio-rad	d.com	Email: antibody sales de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M382404:210513'

Printed on 01 May 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint