

Datasheet: 1720-9007 **BATCH NUMBER 156661**

Description:	GOAT ANTI RAT CALCITONIN GENE-RELATED PEPTIDE
Specificity:	CGRP
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen				
Immunohistology - Paraffin				
Immunohistology - Resin				
ELISA				1/500 - 1/2500
Western Blotting			•	
Immunofluorescence				

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

Target Species	Det
rarget opecies	Rat
Species Cross Reactivity	Reacts with: Mouse, Guinea Pig, Emu N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated fo further information.
Product Form	Purified Ig - liquid
Preparation	Purified Ig prepared by affinity chromatography on Protein G
Buffer Solution	Phosphate buffered saline

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 5.0 mg/ml
Immunogen	Synthetic rat Tyr-CGRP (23-37) conjugated to gamma globulin.
External Database Links	UniProt: P01256 Related reagents Entrez Gene: 24241 Calca Related reagents
Synonyms	Calc
RRID	AB_2290729
Specificity	Goat anti Rat Calcitonin Gene-Related Peptide antibody recognizes Calcitonin gene-related peptide, also known as CGRP, a neuropeptide that acts as a vasodilator and

acids 1-37) and the C-terminal fragment (23-37).

References

1. Collins, J.J. *et al.* (2000) Distribution and origin of secretoneurin-immunoreactive nerves in the female rat uterus. Neuroscience. 95 (1): 255-64.

plays a role in in the pathophysiology of migraine (Recober & Russo 2009). Goat anti Rat Calcitonin Gene-Related Peptide antibody reacts with both the whole molecule (amino

- 2. Pritz, M.B. & Stritzel, M.E. (1988) Thalamic nuclei that project to reptilian telencephalon lack GABA and GAD immunoreactive neurons and puncta. <u>Brain Res. 457 (1): 154-9.</u>
- 3. Pritz, M.B. & Stritzel, M.E. (1989) Reptilian dorsal column nucleus lacks GAD immunoreactive neurons. <u>Brain Res. 503 (1): 175-9.</u>
- 4. Fan, W. *et al.* (2010) Structural and cellular features in metaphyseal and diaphyseal periosteum of osteoporotic rats. J Mol Histol. 41: 51-60.
- 5. Hamed, K. *et al.* (2011) Changes in cutaneous innervation in patients with chronic pain after burns. Burns. 37: 631-7
- 6. Brock, J.A. *et al.* (2007) Postnatal androgen deprivation dissociates the development of smooth muscle innervation from functional neurotransmission in mouse vas deferens. <u>J Physiol. 581: 665-78.</u>
- 7. Gnanamanickam, G.J. and Llewellyn-Smith, I.J. (2011) Innervation of the rat uterus at estrus: a study in full-thickness, immunoperoxidase-stained whole-mount preparations. <u>J Comp Neurol. 519: 621-43.</u>
- 8. Marchant, N.J. *et al.* (2007) Coexpression of prodynorphin and corticotrophin-releasing hormone in the rat central amygdala: evidence of two distinct endogenous opioid systems in the lateral division. <u>J Comp Neurol</u>. 504: 702-15.
- 9. Golden, J.P. *et al.* (2010) RET signaling is required for survival and normal function of nonpeptidergic nociceptors. <u>J Neurosci. 30: 3983-94.</u>
- 10. Ikeda, E. *et al.* (2009) Fully functional bioengineered tooth replacement as an organ replacement therapy. <u>Proc Natl Acad Sci U S A. 106: 13475-80.</u>
- 11. Iliff, J.J. et al. (2010) Epoxyeicosatrienoic acids are endogenous regulators of

- vasoactive neuropeptide release from trigeminal ganglion neurons. <u>J Neurochem. 115:</u> 1530-42.
- 12. Iliff, J.J. *et al.* (2009) Epoxyeicosanoids as mediators of neurogenic vasodilation in cerebral vessels. Am J Physiol Heart Circ Physiol. 296: H1352-63.
- 13. Tague, S.E. and Smith, P.G. (2011) Vitamin D receptor and enzyme expression in dorsal root ganglia of adult female rats: modulation by ovarian hormones. <u>J Chem Neuroanat.</u> 41: 1-12.
- 14. Zou, M. *et al.* (2012) Brn3a/Pou4f1 regulates dorsal root ganglion sensory neuron specification and axonal projection into the spinal cord. <u>Dev Biol.</u> 364: 114-27.
- 15. Chucair-Elliott, A.J. *et al.* (2015) Degeneration and regeneration of corneal nerves in response to HSV-1 infection. Invest Ophthalmol Vis Sci. 56 (2): 1097-107.
- 16. Yu, W.M. *et al.* (2009) Disruption of laminin in the peripheral nervous system impedes nonmyelinating Schwann cell development and impairs nociceptive sensory function. <u>Glia.</u> <u>57: 850-9.</u>
- 17. Zimmerman, A.L. *et al.* (2012) Monoaminergic Modulation of Spinal Viscero-Sympathetic Function in the Neonatal Mouse Thoracic Spinal Cord PLoS One. 7: e47213.
- 18. Drummond, E.S. *et al.* (2014) Increased expression of cutaneous α1-adrenoceptors after chronic constriction injury in rats. <u>J Pain. 15 (2): 188-96.</u>
- 19. Li, Z. *et al.* (2014) Activation of MrgC receptor inhibits N-type calcium channels in small-diameter primary sensory neurons in mice. <u>Pain. 155 (8): 1613-21.</u>
- 20. Drummond, P.D. *et al.* (2014) Upregulation of α 1-adrenoceptors on cutaneous nerve fibres after partial sciatic nerve ligation and in complex regional pain syndrome type II. Pain. 155: 606-16.
- 21. Marvaldi L *et al.* (2015) Enhanced axon outgrowth and improved long-distance axon regeneration in sprouty2 deficient mice. Dev Neurobiol. 75 (3): 217-31.
- 22. Pitzer, C. *et al.* (2008) Granulocyte-colony stimulating factor improves outcome in a mouse model of amyotrophic lateral sclerosis. Brain. 131: 3335-47.
- 23. Chucair-Elliott, A.J. *et al.* (2015) Degeneration and regeneration of corneal nerves in response to HSV-1 infection. Invest Ophthalmol Vis Sci. 56 (2): 1097-107.
- 24. Weir, K.A. and Lunam, C.A. (2006) Immunohistochemical study of cutaneous nerves in the emu. Cell Tissue Res. 326: 697-705.
- 25. Valtcheva, M.V. *et al.* (2015) Enhanced nonpeptidergic intraepidermal fiber density and an expanded subset of chloroquine-responsive trigeminal neurons in a mouse model of dry skin itch. J Pain. 16 (4): 346-56.
- 26. Payne, S.C. *et al.* (2015) Regeneration of sensory but not motor axons following visceral nerve injury. <u>Exp Neurol. 266: 127-42.</u>
- 27. Van Steenwinckel, J. *et al.* (2015) Stromal cell-derived CCL2 drives neuropathic pain states through myeloid cell infiltration in injured nerve. <u>Brain Behav Immun. 45: 198-210.</u>
- 28. Wong, A.W. *et al.* (2015) Neurite outgrowth in normal and injured primary sensory neurons reveals different regulation by nerve growth factor (NGF) and artemin. <u>Mol Cell Neurosci. 65: 125-34.</u>
- 29. Watanabe, M. *et al.* (2015) Expression and Regulation of Cav3.2 T-Type Calcium Channels during Inflammatory Hyperalgesia in Mouse Dorsal Root Ganglion Neurons. <u>PLoS One. 10 (5): e0127572.</u>
- 30. O'Brien, D.E. *et al.* (2015) ERK2 Alone Drives Inflammatory Pain But Cooperates with ERK1 in Sensory Neuron Survival. J Neurosci. 35 (25): 9491-507.
- 31. Sheahan, T.D. et al. (2015) Voluntary Exercise Training: Analysis of Mice in Uninjured,

Inflammatory, and Nerve-Injured Pain States. PLoS One. 10 (7): e0133191.

- 32. Park, S.I. *et al.* (2015) Soft, stretchable, fully implantable miniaturized optoelectronic systems for wireless optogenetics. <u>Nat Biotechnol.</u> 33 (12): 1280-1286.
- 33. Vogelaar CF *et al.* (2015) Pharmacological Suppression of CNS Scarring by Deferoxamine Reduces Lesion Volume and Increases Regeneration in an *In Vitro* Model for Astroglial-Fibrotic Scarring and in Rat Spinal Cord Injury *In Vivo*. <u>PLoS One. 10 (7):</u> <u>e0134371.</u>
- 34. Chetty, R. *et al.* (2006) Pancreatic endocrine tumour with ductules: further observations of an unusual histological subtype. <u>Pathology. 38 (1): 5-9.</u>
- 35. Huang, A.Y. & Wu, S.Y. (2015) Calcitonin Gene-Related Peptide Reduces Taste-Evoked ATP Secretion from Mouse Taste Buds. J Neurosci. 35 (37): 12714-24.
- 36. Lin, S.H. *et al.* (2016) Evidence for the involvement of ASIC3 in sensory mechanotransduction in proprioceptors. Nat Commun. 7: 11460.
- 37. Kim, Y.S. *et al.* (2016) Coupled Activation of Primary Sensory Neurons Contributes to Chronic Pain. <u>Neuron. 91 (5): 1085-96.</u>
- 38. Takahashi, N. *et al.* (2016) Morphology of P2X3-immunoreactive nerve endings in the rat laryngeal mucosa. <u>Histochem Cell Biol. 145 (2): 131-46.</u>
- 39. Takiguchi, M. *et al.* (2018) Neonatal spinal injury induces *de novo* projections of primary afferents to the lumbosacral intermediolateral nucleus in rats. <u>IBRO Rep. 4: 1-6.</u>
- 40. Sun, Y. *et al.* (2020) Somatostatin neurons in the central amygdala mediate anxiety by disinhibition of the central sublenticular extended amygdala. <u>Mol Psychiatry. Oct 01 [Epub ahead of print].</u>
- 41. Rytel, L. & Całka, J. (2020) Aspirin Administration Affects Neurochemical Characterization of Substance P-Like Immunoreactive (SP-LI) Nodose Ganglia Neurons Supplying the Porcine Stomach. Biomed Res Int. 2020: 1049179.

Storage

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted.

Avoid repeated freezing and thawing as this may denature the antibody.

Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/1720-9007 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) FITC, HRP

Recommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

Fax: +1 919 878 3751 Fax: +44 (0)1865 852 739 Fax: +49 (0) 89 8090 95 50 To Email: antibody_sales_us@bio-rad.com Email: antibody_sales_us@bio-rad.com Email: antibody_sales_de@bio-rad.com a

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M362924:200528'

Printed on 21 Mar 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint