# Datasheet: 1720-9007 BATCH NUMBER 160792

Description:	GOAT ANTI RAT CALCITONIN GENE-RELATED PEPTIDE
Specificity:	CGRP
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 ml

#### **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal					
	communications from the originators. Please refer to references indicated for further					
	information. For general	orotocol re	ecommen	dations, please visit <u>v</u>	<u>/ww.bio-</u>	
	rad-antibodies.com/protocols.					
		Yes	No	Not Determined	Suggested Dilution	
	Immunohistology - Frozen					
	Immunohistology - Paraffin	•				
	Immunohistology - Resin					
	ELISA	•			1/500 - 1/2500	
	Western Blotting			•		
	Immunofluorescence					
	Where this product has n	ot been te	ested for	use in a particular tecl	nnique this does not	
	necessarily exclude its use in such procedures. Suggested working dilutions are given as					
	a guide only. It is recomm	nended the	at the use	er titrates the product	for use in their own	
	system using the appropriate negative/positive controls.					
Target Species	Rat					
Species Cross	Reacts with: Mouse, Guir	nea Pig, E	mu			
Reactivity	<b>N.B.</b> Antibody reactivity a	nd workin	ng conditi	ons may vary betweer	n species. Cross	
	reactivity is derived from	testing wit	thin our la	aboratories, peer-revie	ewed publications or	
	personal communications	s from the	originato	rs. Please refer to refe	erences indicated for	
	further information.					
Product Form	Purified la - liquid					
	i annoa igʻilqala					
Preparation	Purified Ig prepared by a	ffinity chro	omatogra	ohy on Protein G		
Buffer Solution	Phosphate buffered salin	e				

Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> )		
Approx. Protein Concentrations	IgG concentration 5.0 mg/ml		
Immunogen	Synthetic rat Tyr-CGRP (23-37) conjugated to gamma globulin.		
External Database Links	UniProt: <u>P01256</u> <u>Related reagents</u>		
	<u>24241</u> Calca <u>Related reagents</u>		
Synonyms	Calc		
RRID	AB_2290729		
Specificity	<b>Goat anti Rat Calcitonin Gene-Related Peptide antibody</b> recognizes Calcitonin gene-related peptide, also known as CGRP, a neuropeptide that acts as a vasodilator and plays a role in in the pathophysiology of migraine ( <u>Recober &amp; Russo 2009</u> ). Goat anti Rat Calcitonin Gene-Related Peptide antibody reacts with both the whole molecule (amino acids 1-37) and the C-terminal fragment (23-37).		
References	<ol> <li>Collins, J.J. <i>et al.</i> (2000) Distribution and origin of secretoneurin-immunoreactive nerves in the female rat uterus. Neuroscience. 95 (1): 255-64.</li> <li>Chetty, R. <i>et al.</i> (2006) Pancreatic endocrine tumour with ductules: further observations of an unusual histological subtype. Pathology. 38 (1): 5-9.</li> <li>Weir, K.A. and Lunam, C.A. (2006) Immunohistochemical study of cutaneous nerves in the emu. <u>Cell Tissue Res. 326: 697-705.</u></li> <li>Brock, J.A. <i>et al.</i> (2007) Postnatal androgen deprivation dissociates the development of smooth muscle innervation from functional neurotransmission in mouse vas deferens. J <u>Physiol. 581: 665-78.</u></li> <li>Marchant, N.J. <i>et al.</i> (2007) Coexpression of prodynorphin and corticotrophin-releasing hormone in the rat central amygdala: evidence of two distinct endogenous opioid systems in the lateral division. J Comp Neurol. 504: 702-15.</li> <li>Pitzer, C. <i>et al.</i> (2008) Granulocyte-colony stimulating factor improves outcome in a mouse model of amyotrophic lateral sclerosis. <u>Brain. 131: 3335-47.</u></li> <li>Ikeda, E. <i>et al.</i> (2009) Fully functional bioengineered tooth replacement as an organ replacement therapy. <u>Proc Natl Acad Sci U S A. 106: 13475-80.</u></li> <li>Iliff, J.J. <i>et al.</i> (2009) Disruption of laminin in the peripheral nervous system impedes nonmyelinating Schwann cell development and impairs nociceptive sensory function. <u>Glia.</u> <u>57: 850-9.</u></li> <li>Golden, J.P. <i>et al.</i> (2010) RET signaling is required for survival and normal function of nonpeptidergic nociceptors. <u>J Neurosci. 30: 3983-94.</u></li> <li>Iliff, J.J. <i>et al.</i> (2010) Epoxyeicosatrienoic acids are endogenous regulators of</li> </ol>		

vasoactive neuropeptide release from trigeminal ganglion neurons. <u>J Neurochem. 115:</u> 1530-42.

12. Fan, W. *et al.* (2010) Structural and cellular features in metaphyseal and diaphyseal periosteum of osteoporotic rats. J Mol Histol. 41: 51-60.

13. Gnanamanickam, G.J. and Llewellyn-Smith, I.J. (2011) Innervation of the rat uterus at estrus: a study in full-thickness, immunoperoxidase-stained whole-mount preparations. J Comp Neurol. 519: 621-43.

14. Tague, S.E. and Smith, P.G. (2011) Vitamin D receptor and enzyme expression in dorsal root ganglia of adult female rats: modulation by ovarian hormones. <u>J Chem</u> <u>Neuroanat. 41: 1-12.</u>

15. Hamed, K. *et al.* (2011) Changes in cutaneous innervation in patients with chronic pain after burns. <u>Burns. 37: 631-7</u>

16. Zou, M. *et al.* (2012) Brn3a/Pou4f1 regulates dorsal root ganglion sensory neuron specification and axonal projection into the spinal cord. <u>Dev Biol. 364: 114-27.</u>

17. Zimmerman, A.L. et al. (2012) Monoaminergic Modulation of Spinal Viscero-

Sympathetic Function in the Neonatal Mouse Thoracic Spinal Cord PLoS One. 7: e47213.

18. Drummond, E.S. *et al.* (2014) Increased expression of cutaneous α1-adrenoceptors after chronic constriction injury in rats. <u>J Pain. 15 (2): 188-96.</u>

19. Drummond, P.D. *et al.* (2014) Upregulation of  $\alpha$ 1-adrenoceptors on cutaneous nerve fibres after partial sciatic nerve ligation and in complex regional pain syndrome type II. <u>Pain. 155: 606-16.</u>

20. Li, Z. *et al.* (2014) Activation of MrgC receptor inhibits N-type calcium channels in small-diameter primary sensory neurons in mice. <u>Pain. 155 (8): 1613-21.</u>

21. Park, S.I. *et al.* (2015) Soft, stretchable, fully implantable miniaturized optoelectronic systems for wireless optogenetics. <u>Nat Biotechnol. 33 (12): 1280-1286.</u>

22. O'Brien, D.E. *et al.* (2015) ERK2 Alone Drives Inflammatory Pain But Cooperates with ERK1 in Sensory Neuron Survival. <u>J Neurosci. 35 (25): 9491-507.</u>

23. Vogelaar CF *et al.* (2015) Pharmacological Suppression of CNS Scarring by Deferoxamine Reduces Lesion Volume and Increases Regeneration in an *In Vitro* Model for Astroglial-Fibrotic Scarring and in Rat Spinal Cord Injury *In Vivo*. <u>PLoS One. 10 (7):</u> <u>e0134371.</u>

24. Sheahan, T.D. *et al.* (2015) Voluntary Exercise Training: Analysis of Mice in Uninjured, Inflammatory, and Nerve-Injured Pain States. <u>PLoS One. 10 (7): e0133191.</u>

25. Huang, A.Y. & Wu, S.Y. (2015) Calcitonin Gene-Related Peptide Reduces Taste-Evoked ATP Secretion from Mouse Taste Buds. <u>J Neurosci. 35 (37): 12714-24.</u>

26. Watanabe, M. *et al.* (2015) Expression and Regulation of Cav3.2 T-Type Calcium Channels during Inflammatory Hyperalgesia in Mouse Dorsal Root Ganglion Neurons. <u>PLoS One. 10 (5): e0127572.</u>

27. Chucair-Elliott, A.J. *et al.* (2015) Degeneration and regeneration of corneal nerves in response to HSV-1 infection. Invest Ophthalmol Vis Sci. 56 (2): 1097-107.

Van Steenwinckel, J. *et al.* (2015) Stromal cell-derived CCL2 drives neuropathic pain states through myeloid cell infiltration in injured nerve. <u>Brain Behav Immun. 45: 198-210.</u>
 Payne, S.C. *et al.* (2015) Regeneration of sensory but not motor axons following visceral nerve injury. <u>Exp Neurol. 266: 127-42.</u>

30. Valtcheva, M.V. *et al.* (2015) Enhanced nonpeptidergic intraepidermal fiber density and an expanded subset of chloroquine-responsive trigeminal neurons in a mouse model of dry skin itch. J Pain. 16 (4): 346-56.

	31. Chucair-Elliott, A.J. et al. (2015) Degeneration and regener	ation of corneal nerves in
	response to HSV-1 infection. Invest Ophthalmol Vis Sci. 56 (2):	<u>1097-107.</u>
	32. Marvaldi L et al. (2015) Enhanced axon outgrowth and impl	roved long-distance axon
	regeneration in sprouty2 deficient mice. Dev Neurobiol. 75 (3):	<u>217-31.</u>
	33. Wong, A.W. et al. (2015) Neurite outgrowth in normal and in	njured primary sensory
	neurons reveals different regulation by nerve growth factor (NG	F) and artemin. <u>Mol Cell</u>
	Neurosci. 65: 125-34.	
	34. Lin, S.H. et al. (2016) Evidence for the involvement of ASIC	C3 in sensory
	mechanotransduction in proprioceptors. Nat Commun. 7: 11460	<u>).</u>
	35. Kim, Y.S. et al. (2016) Coupled Activation of Primary Senso	ory Neurons Contributes to
	Chronic Pain. <u>Neuron. 91 (5): 1085-96.</u>	
	36. Takahashi, N. et al. (2016) Morphology of P2X3-immunorea	active nerve endings in the
	rat laryngeal mucosa. <u>Histochem Cell Biol. 145 (2): 131-46.</u>	
	37. Takiguchi, M. et al. (2018) Neonatal spinal injury induces de	e <i>novo</i> projections of
	primary afferents to the lumbosacral intermediolateral nucleus i	n rats. <u>IBRO Rep. 4: 1-6.</u>
	38. Sun, Y. et al. (2020) Somatostatin neurons in the central an	nygdala mediate anxiety by
	disinhibition of the central sublenticular extended amygdala. Mo	ol Psychiatry. Oct 01 [Epub
	ahead of print].	
	39. Rytel, L. & Całka, J. (2020) Aspirin Administration Affects N	eurochemical
	Characterization of Substance P-Like Immunoreactive (SP-LI) N	Nodose Ganglia Neurons
	Supplying the Porcine Stomach. Biomed Res Int. 2020: 104917	<u>′9.</u>
	40. Morellini, N. et al. (2020) Decreased neural expression of the	ne noradrenaline
	transporter in the papillary dermis after partial sciatic nerve lesi	on. <u>J Chem Neuroanat.</u>
	<u>107: 101806.</u>	
	41. Lin, J.H. et al. (2021) ATF3-Expressing Large-Diameter Ser	nsory Afferents at Acute
	Stage as Bio-Signatures of Persistent Pain Associated with Lur	nbar Radiculopathy. <u>Cells.</u>
	<u>10 (5): 992</u>	
Storage	This product is shipped at ambient temperature. It is recommer	nded to aliquot and store at
	-20°C on receipt. When thawed, aliquot the sample as needed.	Keep aliquots at 2-8°C for
	short term use (up to 4 weeks) and store the remaining aliquots	s at -20°C.
	Avoid repeated freezing and thawing as this may denature the	antibody. Storage in
	frost-free freezers is not recommended.	
Guarantee	12 months from date of despatch	
	,	
Health And Safety	Material Safety Datasheet documentation #10040 available at:	
Information	https://www.bio-rad-antibodies.com/SDS/1720-9007	
	10040	
Regulatory	For research purposes only	
······································		

# **Related Products**

# **Recommended Secondary Antibodies**

Rabbit Anti Goat IgG (Fc) (STAR122...) FITC, HRP

#### ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio	-rad.com	Email: antibody_sales_uk@bio-rac	l.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M381311:210512'

#### Printed on 21 Mar 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint