

Datasheet: 1720-9007

BATCH NUMBER 161056

Description:	GOAT ANTI RAT CALCITONIN GENE-RELATED PEPTIDE
Specificity:	CGRP
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	■			
Immunohistology - Paraffin	■			
Immunohistology - Resin		■		
ELISA	■			1/500 - 1/2500
Western Blotting			■	
Immunofluorescence	■			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

Target Species

Rat

Species Cross Reactivity

Reacts with: Mouse, Guinea Pig, Emu

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified Ig - liquid

Preparation

Purified Ig prepared by affinity chromatography on Protein G

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 5.0 mg/ml
Immunogen	Synthetic rat Tyr-CGRP (23-37) conjugated to gamma globulin.
External Database Links	<p>UniProt: P01256 Related reagents</p> <p>Entrez Gene: 24241 Calca Related reagents</p>
Synonyms	Calc
RRID	AB_2290729
Specificity	Goat anti Rat Calcitonin Gene-Related Peptide antibody recognizes Calcitonin gene-related peptide, also known as CGRP, a neuropeptide that acts as a vasodilator and plays a role in in the pathophysiology of migraine (Recober & Russo 2009). Goat anti Rat Calcitonin Gene-Related Peptide antibody reacts with both the whole molecule (amino acids 1-37) and the C-terminal fragment (23-37).
References	<ol style="list-style-type: none"> Collins, J.J. <i>et al.</i> (2000) Distribution and origin of secretoneurin-immunoreactive nerves in the female rat uterus. Neuroscience. 95 (1): 255-64. Chetty, R. <i>et al.</i> (2006) Pancreatic endocrine tumour with ductules: further observations of an unusual histological subtype. Pathology. 38 (1): 5-9. Weir, K.A. and Lunam, C.A. (2006) Immunohistochemical study of cutaneous nerves in the emu. Cell Tissue Res. 326: 697-705. Brock, J.A. <i>et al.</i> (2007) Postnatal androgen deprivation dissociates the development of smooth muscle innervation from functional neurotransmission in mouse vas deferens. J Physiol. 581: 665-78. Marchant, N.J. <i>et al.</i> (2007) Coexpression of prodynorphin and corticotrophin-releasing hormone in the rat central amygdala: evidence of two distinct endogenous opioid systems in the lateral division. J Comp Neurol. 504: 702-15. Pitzer, C. <i>et al.</i> (2008) Granulocyte-colony stimulating factor improves outcome in a mouse model of amyotrophic lateral sclerosis. Brain. 131: 3335-47. Ikeda, E. <i>et al.</i> (2009) Fully functional bioengineered tooth replacement as an organ replacement therapy. Proc Natl Acad Sci U S A. 106: 13475-80. Iliff, J.J. <i>et al.</i> (2009) Epoxyeicosanoids as mediators of neurogenic vasodilation in cerebral vessels. Am J Physiol Heart Circ Physiol. 296: H1352-63. Yu, W.M. <i>et al.</i> (2009) Disruption of laminin in the peripheral nervous system impedes nonmyelinating Schwann cell development and impairs nociceptive sensory function. Glia. 57: 850-9. Golden, J.P. <i>et al.</i> (2010) RET signaling is required for survival and normal function of nonpeptidergic nociceptors. J Neurosci. 30: 3983-94. Iliff, J.J. <i>et al.</i> (2010) Epoxyeicosatrienoic acids are endogenous regulators of

- vasoactive neuropeptide release from trigeminal ganglion neurons. [J Neurochem. 115: 1530-42.](#)
12. Fan, W. *et al.* (2010) Structural and cellular features in metaphyseal and diaphyseal periosteum of osteoporotic rats. [J Mol Histol. 41: 51-60.](#)
 13. Gnanamanickam, G.J. and Llewellyn-Smith, I.J. (2011) Innervation of the rat uterus at estrus: a study in full-thickness, immunoperoxidase-stained whole-mount preparations. [J Comp Neurol. 519: 621-43.](#)
 14. Tague, S.E. and Smith, P.G. (2011) Vitamin D receptor and enzyme expression in dorsal root ganglia of adult female rats: modulation by ovarian hormones. [J Chem Neuroanat. 41: 1-12.](#)
 15. Hamed, K. *et al.* (2011) Changes in cutaneous innervation in patients with chronic pain after burns. [Burns. 37: 631-7](#)
 16. Zou, M. *et al.* (2012) Brn3a/Pou4f1 regulates dorsal root ganglion sensory neuron specification and axonal projection into the spinal cord. [Dev Biol. 364: 114-27.](#)
 17. Zimmerman, A.L. *et al.* (2012) Monoaminergic Modulation of Spinal Viscero-Sympathetic Function in the Neonatal Mouse Thoracic Spinal Cord [PLoS One. 7: e47213.](#)
 18. Drummond, E.S. *et al.* (2014) Increased expression of cutaneous α 1-adrenoceptors after chronic constriction injury in rats. [J Pain. 15 \(2\): 188-96.](#)
 19. Drummond, P.D. *et al.* (2014) Upregulation of α 1-adrenoceptors on cutaneous nerve fibres after partial sciatic nerve ligation and in complex regional pain syndrome type II. [Pain. 155: 606-16.](#)
 20. Li, Z. *et al.* (2014) Activation of MrgC receptor inhibits N-type calcium channels in small-diameter primary sensory neurons in mice. [Pain. 155 \(8\): 1613-21.](#)
 21. Park, S.I. *et al.* (2015) Soft, stretchable, fully implantable miniaturized optoelectronic systems for wireless optogenetics. [Nat Biotechnol. 33 \(12\): 1280-1286.](#)
 22. O'Brien, D.E. *et al.* (2015) ERK2 Alone Drives Inflammatory Pain But Cooperates with ERK1 in Sensory Neuron Survival. [J Neurosci. 35 \(25\): 9491-507.](#)
 23. Vogelaar CF *et al.* (2015) Pharmacological Suppression of CNS Scarring by Deferoxamine Reduces Lesion Volume and Increases Regeneration in an *In Vitro* Model for Astroglial-Fibrotic Scarring and in Rat Spinal Cord Injury *In Vivo*. [PLoS One. 10 \(7\): e0134371.](#)
 24. Sheahan, T.D. *et al.* (2015) Voluntary Exercise Training: Analysis of Mice in Uninjured, Inflammatory, and Nerve-Injured Pain States. [PLoS One. 10 \(7\): e0133191.](#)
 25. Huang, A.Y. & Wu, S.Y. (2015) Calcitonin Gene-Related Peptide Reduces Taste-Evoked ATP Secretion from Mouse Taste Buds. [J Neurosci. 35 \(37\): 12714-24.](#)
 26. Watanabe, M. *et al.* (2015) Expression and Regulation of Cav3.2 T-Type Calcium Channels during Inflammatory Hyperalgesia in Mouse Dorsal Root Ganglion Neurons. [PLoS One. 10 \(5\): e0127572.](#)
 27. Chucair-Elliott, A.J. *et al.* (2015) Degeneration and regeneration of corneal nerves in response to HSV-1 infection. [Invest Ophthalmol Vis Sci. 56 \(2\): 1097-107.](#)
 28. Van Steenwinckel, J. *et al.* (2015) Stromal cell-derived CCL2 drives neuropathic pain states through myeloid cell infiltration in injured nerve. [Brain Behav Immun. 45: 198-210.](#)
 29. Payne, S.C. *et al.* (2015) Regeneration of sensory but not motor axons following visceral nerve injury. [Exp Neurol. 266: 127-42.](#)
 30. Valtcheva, M.V. *et al.* (2015) Enhanced nonpeptidergic intraepidermal fiber density and an expanded subset of chloroquine-responsive trigeminal neurons in a mouse model of dry skin itch. [J Pain. 16 \(4\): 346-56.](#)

31. Chucair-Elliott, A.J. *et al.* (2015) Degeneration and regeneration of corneal nerves in response to HSV-1 infection. [Invest Ophthalmol Vis Sci. 56 \(2\): 1097-107.](#)
32. Marvaldi L *et al.* (2015) Enhanced axon outgrowth and improved long-distance axon regeneration in sprouty2 deficient mice. [Dev Neurobiol. 75 \(3\): 217-31.](#)
33. Wong, A.W. *et al.* (2015) Neurite outgrowth in normal and injured primary sensory neurons reveals different regulation by nerve growth factor (NGF) and artemin. [Mol Cell Neurosci. 65: 125-34.](#)
34. Lin, S.H. *et al.* (2016) Evidence for the involvement of ASIC3 in sensory mechanotransduction in proprioceptors. [Nat Commun. 7: 11460.](#)
35. Kim, Y.S. *et al.* (2016) Coupled Activation of Primary Sensory Neurons Contributes to Chronic Pain. [Neuron. 91 \(5\): 1085-96.](#)
36. Takahashi, N. *et al.* (2016) Morphology of P2X3-immunoreactive nerve endings in the rat laryngeal mucosa. [Histochem Cell Biol. 145 \(2\): 131-46.](#)
37. Takiguchi, M. *et al.* (2018) Neonatal spinal injury induces *de novo* projections of primary afferents to the lumbosacral intermediolateral nucleus in rats. [IBRO Rep. 4: 1-6.](#)
38. Sun, Y. *et al.* (2020) Somatostatin neurons in the central amygdala mediate anxiety by disinhibition of the central subnucleus extended amygdala. [Mol Psychiatry. Oct 01 \[Epub ahead of print\].](#)
39. Rytel, L. & Calka, J. (2020) Aspirin Administration Affects Neurochemical Characterization of Substance P-Like Immunoreactive (SP-LI) Nodose Ganglia Neurons Supplying the Porcine Stomach. [Biomed Res Int. 2020: 1049179.](#)
40. Morellini, N. *et al.* (2020) Decreased neural expression of the noradrenaline transporter in the papillary dermis after partial sciatic nerve lesion. [J Chem Neuroanat. 107: 101806.](#)
41. Lin, J.H. *et al.* (2021) ATF3-Expressing Large-Diameter Sensory Afferents at Acute Stage as Bio-Signatures of Persistent Pain Associated with Lumbar Radiculopathy. [Cells. 10 \(5\): 992](#)

Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/1720-9007 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) [FITC](#), [HRP](#)

Recommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M381311:210512'

Printed on 21 Mar 2025

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)