

Datasheet: 1720-9007 **BATCH NUMBER 161057**

Description:	GOAT ANTI RAT CALCITONIN GENE-RELATED PEPTIDE
Specificity:	CGRP
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	•			
Immunohistology - Paraffin	•			
Immunohistology - Resin		•		
ELISA	•			1/500 - 1/2500
Western Blotting				
Immunofluorescence	•			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

Target Species	Rat	
Species Cross Reactivity	Reacts with: Mouse, Guinea Pig, Emu N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.	
Product Form	Purified Ig - liquid	
Preparation	Purified Ig prepared by affinity chromatography on Protein G	
Buffer Solution	Phosphate buffered saline	

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 5.0 mg/ml
Immunogen	Synthetic rat Tyr-CGRP (23-37) conjugated to gamma globulin.
External Database Links	UniProt: P01256 Related reagents
	Entrez Gene: 24241 Calca Related reagents
Synonyms	Calc
RRID	AB_2290729
Specificity	Goat anti Rat Calcitonin Gene-Related Peptide antibody recognizes Calcitonin gene-related peptide, also known as CGRP, a neuropeptide that acts as a vasodilator and plays a role in in the pathophysiology of migraine (Recober & Russo 2009). Goat anti Rat

acids 1-37) and the C-terminal fragment (23-37).

References

1. Collins, J.J. *et al.* (2000) Distribution and origin of secretoneurin-immunoreactive nerves in the female rat uterus. Neuroscience. 95 (1): 255-64.

Calcitonin Gene-Related Peptide antibody reacts with both the whole molecule (amino

- 2. Chetty, R. *et al.* (2006) Pancreatic endocrine tumour with ductules: further observations of an unusual histological subtype. <u>Pathology. 38 (1): 5-9.</u>
- 3. Weir, K.A. and Lunam, C.A. (2006) Immunohistochemical study of cutaneous nerves in the emu. <u>Cell Tissue Res. 326: 697-705.</u>
- 4. Brock, J.A. *et al.* (2007) Postnatal androgen deprivation dissociates the development of smooth muscle innervation from functional neurotransmission in mouse vas deferens. <u>J Physiol. 581: 665-78.</u>
- 5. Marchant, N.J. *et al.* (2007) Coexpression of prodynorphin and corticotrophin-releasing hormone in the rat central amygdala: evidence of two distinct endogenous opioid systems in the lateral division. <u>J Comp Neurol</u>. 504: 702-15.
- 6. Pitzer, C. *et al.* (2008) Granulocyte-colony stimulating factor improves outcome in a mouse model of amyotrophic lateral sclerosis. <u>Brain. 131: 3335-47.</u>
- 7. lkeda, E. *et al.* (2009) Fully functional bioengineered tooth replacement as an organ replacement therapy. <u>Proc Natl Acad Sci U S A. 106: 13475-80.</u>
- 8. Iliff, J.J. *et al.* (2009) Epoxyeicosanoids as mediators of neurogenic vasodilation in cerebral vessels. <u>Am J Physiol Heart Circ Physiol.</u> 296: H1352-63.
- 9. Yu, W.M. *et al.* (2009) Disruption of laminin in the peripheral nervous system impedes nonmyelinating Schwann cell development and impairs nociceptive sensory function. <u>Glia.</u> 57: 850-9.
- 10. Golden, J.P. *et al.* (2010) RET signaling is required for survival and normal function of nonpeptidergic nociceptors. <u>J Neurosci. 30: 3983-94.</u>
- 11. Iliff, J.J. et al. (2010) Epoxyeicosatrienoic acids are endogenous regulators of

- vasoactive neuropeptide release from trigeminal ganglion neurons. <u>J Neurochem. 115:</u> 1530-42.
- 12. Fan, W. *et al.* (2010) Structural and cellular features in metaphyseal and diaphyseal periosteum of osteoporotic rats. <u>J Mol Histol. 41: 51-60.</u>
- 13. Gnanamanickam, G.J. and Llewellyn-Smith, I.J. (2011) Innervation of the rat uterus at estrus: a study in full-thickness, immunoperoxidase-stained whole-mount preparations. <u>J</u> Comp Neurol. 519: 621-43.
- 14. Tague, S.E. and Smith, P.G. (2011) Vitamin D receptor and enzyme expression in dorsal root ganglia of adult female rats: modulation by ovarian hormones. <u>J Chem Neuroanat. 41: 1-12.</u>
- 15. Hamed, K. *et al.* (2011) Changes in cutaneous innervation in patients with chronic pain after burns. <u>Burns. 37: 631-7</u>
- 16. Zou, M. *et al.* (2012) Brn3a/Pou4f1 regulates dorsal root ganglion sensory neuron specification and axonal projection into the spinal cord. Dev Biol. 364: 114-27.
- 17. Zimmerman, A.L. *et al.* (2012) Monoaminergic Modulation of Spinal Viscero-Sympathetic Function in the Neonatal Mouse Thoracic Spinal Cord PLoS One. 7: e47213.
- 18. Drummond, E.S. *et al.* (2014) Increased expression of cutaneous α1-adrenoceptors after chronic constriction injury in rats. <u>J Pain. 15 (2): 188-96.</u>
- 19. Drummond, P.D. *et al.* (2014) Upregulation of α 1-adrenoceptors on cutaneous nerve fibres after partial sciatic nerve ligation and in complex regional pain syndrome type II. Pain. 155: 606-16.
- 20. Li, Z. *et al.* (2014) Activation of MrgC receptor inhibits N-type calcium channels in small-diameter primary sensory neurons in mice. Pain. 155 (8): 1613-21.
- 21. Park, S.I. *et al.* (2015) Soft, stretchable, fully implantable miniaturized optoelectronic systems for wireless optogenetics. Nat Biotechnol. 33 (12): 1280-1286.
- 22. O'Brien, D.E. *et al.* (2015) ERK2 Alone Drives Inflammatory Pain But Cooperates with ERK1 in Sensory Neuron Survival. J Neurosci. 35 (25): 9491-507.
- 23. Vogelaar CF *et al.* (2015) Pharmacological Suppression of CNS Scarring by Deferoxamine Reduces Lesion Volume and Increases Regeneration in an *In Vitro* Model for Astroglial-Fibrotic Scarring and in Rat Spinal Cord Injury *In Vivo*. <u>PLoS One. 10 (7):</u> e0134371.
- 24. Sheahan, T.D. *et al.* (2015) Voluntary Exercise Training: Analysis of Mice in Uninjured, Inflammatory, and Nerve-Injured Pain States. <u>PLoS One. 10 (7): e0133191.</u>
- 25. Huang, A.Y. & Wu, S.Y. (2015) Calcitonin Gene-Related Peptide Reduces Taste-Evoked ATP Secretion from Mouse Taste Buds. J Neurosci. 35 (37): 12714-24.
- 26. Watanabe, M. *et al.* (2015) Expression and Regulation of Cav3.2 T-Type Calcium Channels during Inflammatory Hyperalgesia in Mouse Dorsal Root Ganglion Neurons. <u>PLoS One. 10 (5): e0127572.</u>
- 27. Chucair-Elliott, A.J. *et al.* (2015) Degeneration and regeneration of corneal nerves in response to HSV-1 infection. <u>Invest Ophthalmol Vis Sci. 56 (2): 1097-107.</u>
- 28. Van Steenwinckel, J. *et al.* (2015) Stromal cell-derived CCL2 drives neuropathic pain states through myeloid cell infiltration in injured nerve. <u>Brain Behav Immun. 45: 198-210.</u>
- 29. Payne, S.C. *et al.* (2015) Regeneration of sensory but not motor axons following visceral nerve injury. <u>Exp Neurol. 266: 127-42.</u>
- 30. Valtcheva, M.V. *et al.* (2015) Enhanced nonpeptidergic intraepidermal fiber density and an expanded subset of chloroquine-responsive trigeminal neurons in a mouse model of dry skin itch. <u>J Pain. 16 (4): 346-56.</u>

- 31. Chucair-Elliott, A.J. *et al.* (2015) Degeneration and regeneration of corneal nerves in response to HSV-1 infection. Invest Ophthalmol Vis Sci. 56 (2): 1097-107.
- 32. Marvaldi L *et al.* (2015) Enhanced axon outgrowth and improved long-distance axon regeneration in sprouty2 deficient mice. <u>Dev Neurobiol. 75 (3): 217-31.</u>
- 33. Wong, A.W. *et al.* (2015) Neurite outgrowth in normal and injured primary sensory neurons reveals different regulation by nerve growth factor (NGF) and artemin. <u>Mol Cell Neurosci. 65: 125-34.</u>
- 34. Lin, S.H. *et al.* (2016) Evidence for the involvement of ASIC3 in sensory mechanotransduction in proprioceptors. <u>Nat Commun. 7: 11460.</u>
- 35. Kim, Y.S. *et al.* (2016) Coupled Activation of Primary Sensory Neurons Contributes to Chronic Pain. Neuron. 91 (5): 1085-96.
- 36. Takahashi, N. *et al.* (2016) Morphology of P2X3-immunoreactive nerve endings in the rat laryngeal mucosa. <u>Histochem Cell Biol. 145 (2): 131-46.</u>
- 37. Takiguchi, M. *et al.* (2018) Neonatal spinal injury induces *de novo* projections of primary afferents to the lumbosacral intermediolateral nucleus in rats. IBRO Rep. 4: 1-6.
- 38. Sun, Y. *et al.* (2020) Somatostatin neurons in the central amygdala mediate anxiety by disinhibition of the central sublenticular extended amygdala. <u>Mol Psychiatry. Oct 01 [Epub ahead of print].</u>
- 39. Rytel, L. & Całka, J. (2020) Aspirin Administration Affects Neurochemical Characterization of Substance P-Like Immunoreactive (SP-LI) Nodose Ganglia Neurons Supplying the Porcine Stomach. <u>Biomed Res Int. 2020: 1049179.</u>
- 40. Morellini, N. *et al.* (2020) Decreased neural expression of the noradrenaline transporter in the papillary dermis after partial sciatic nerve lesion. <u>J Chem Neuroanat.</u> 107: 101806.
- 41. Lin, J.H. *et al.* (2021) ATF3-Expressing Large-Diameter Sensory Afferents at Acute Stage as Bio-Signatures of Persistent Pain Associated with Lumbar Radiculopathy. <u>Cells.</u> 10 (5): 992

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/1720-9007 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) FITC, HRP

Recommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M381311:210512'

Printed on 21 Mar 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint