

Datasheet: 1940-0004 **BATCH NUMBER 171273** 

Description:	SHEEP ANTI HUMAN CERULOPLASMIN
Specificity:	CERULOPLASMIN
Format:	Purified
<b>Product Type:</b>	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 ml

## **Product Details**

#### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="https://www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	•			
Immunohistology - Paraffin				
ELISA	-			
Western Blotting			•	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN <sub>3</sub> )
Approx. Protein Concentrations	IgG concentration 5.0 mg/ml
Immunogen	Highly pure human ceruloplasmin.

External Database Links	UniProt:		
LINKS	P00450 Related reagents		
	Entrez Gene:		
	1356 CP Related reagents		
RRID	AB_2085556		
Specificity	Sheep anti Human ceruloplasmin antibody detects ceruloplasmin, a ~132kDa metalloprotein that is secreted into the plasma by liver hepatocytes and activated monocytes/macrophages. Ceruloplasmin functions in copper transport, iron homeostasis, antioxidant defence, angiogenesis and coagulation.		
	Deficiency in ceruloplasmin is termed aceruloplasminemia, an autosomal recessive disorder which leads to iron accumulation and tissue damage, in particular in the brain and visceral organs. Clinical features include retinal degeneration, diabetes mellitus and neurological disturbances.		
	Elevated plasma ceruloplasmin has been shown to pose an increased risk in atherosclerosis and myocardial infarction.		
ELISA	This antibody has been used at dilutions as high as 1/400K in a simple ELISA with 0.1 $\mu$ g of antigen coated per well.		
References	1. Balmaña, M. <i>et al.</i> (2015) Identification of potential pancreatic cancer serum markers: Increased sialyl-Lewis X on ceruloplasmin. Clin Chim Acta. 442: 56-62.		
Further Reading	1. Mazumder, B. <i>et al.</i> (2005) Regulation of macrophage ceruloplasmin gene expression: one paradigm of 3'-UTR-mediated translational control. <u>Mol Cells. 20 (2): 167-72.</u>		
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.		
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.		
Guarantee	12 months from date of despatch		
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/1940-0004">https://www.bio-rad-antibodies.com/SDS/1940-0004</a> 10040		
Regulatory	For research purposes only		

# Related Products

# **Recommended Secondary Antibodies**

Rabbit Anti Sheep IgG (H/L) (5184-2304...) Biotin

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M427914:240301'

### Printed on 21 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint