

Datasheet: 2100-0657

Description:	MOUSE ANTI HUMAN CHYMOTRYPSIN
Specificity:	CHYMOTRYPSIN
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	CHYMO10-104.5 (4E1)
Isotype:	lgG3
Quantity:	0.2 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	•			
Immunohistology - Paraffin	-			
ELISA	-			
Western Blotting	-			
Immunofluorescence	-			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

Target Species	Human
Species Cross Reactivity	Reacts with: Mouse, Rat N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications personal communications from the originators. Please refer to references indicated further information.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G
Buffer Solution	Phosphate buffered saline
Preservative	<0.1% Sodium Azide (NaN ₃)

Stabilisers

Approx. Protein Concentrations	IgG concentration 1.0 mg/ml				
Immunogen	Purified human pancreatic chymotrypsin.				
External Database Links	UniProt: Q99895 Related reagents				
	Entrez Gene: 11330 CTRC Related reagents				
Synonyms	CLCR				
RRID	AB_2261171				
Specificity	Mouse anti Human chymotrypsin antibody, clone 4E1 recognizes human chymotrypsin-C, also known as chymotrypsin or caldecrin. Chymotrypsin is a 239 amino acid ~30 kDa protease with an additional 13 amino acid propeptide region and a 16 amino acid signal peptide.				
	Variations in the CTRC gene has been associated with susceptibility to hereditary, pancreatitis (<u>PCTT</u>), a disease characterized by pancreatic inflammation and destruction of the parenchyma (<u>Beer et al. 2013</u>).				
References	1. Jimenez, R.E. <i>et al.</i> (1999) Immunohistochemical characterization of pancreatic tumors				

- nors induced by dimethylbenzanthracene in rats. Am J Pathol. 154 (4): 1223-9.
- 2. Larina, O.et al. (2007) Dynamic regulation of the large exocytotic fusion pore in pancreatic acinar cells. Mol Biol Cell. 18:3502-11.
- 3. Bockman, D.E. et al. (2003) Origin and development of the precursor lesions in experimental pancreatic cancer in rats. Lab Invest. 83 (6): 853-9.
- 4. Li, H. et al. (2009) The Ink4/Arf locus is a barrier for iPS cell reprogramming. Nature. 460: 1136-9.
- 5. Vincent, D.F. et al. (2009) Inactivation of TIF1gamma cooperates with Kras to induce cystic tumors of the pancreas. PLoS Genet. 5: e1000575.
- 6. Behrendorff, N. et al. (2011) Vesicle-associated membrane protein 8 (VAMP8) is a SNARE (soluble N-ethylmaleimide-sensitive factor attachment protein receptor) selectively required for sequential granule-to-granule fusion. J Biol Chem. 286 (34): 29627-34.
- 7. Guerra, C. et al. (2011) Pancreatitis-induced inflammation contributes to pancreatic cancer by inhibiting oncogene-induced senescence. Cancer Cell. 19: 728-39.
- 8. Kanayama, K. et al. (2016) Cytological findings of an ectopic pancreas of the stomach obtained at endoscopic ultrasound-guided fine needle aspiration, differential diagnosis from acinar cell carcinoma: a case report. Cytopathology. 27 (5): 379-81.
- 9. Vincent DF et al. (2012) Tif1γ suppresses murine pancreatic tumoral transformation by a Smad4-independent pathway. Am J Pathol. 180 (6): 2214-21.
- 10. Kato, Y. et al. (2014) Ectopic tissue consisting of a mixture of glandular gastric,

intestinal, and exocrine pancreatic tissue in the forestomach of a rat. <u>J Toxicol Pathol. 27</u> (1): 87-90.

11. Bansal, S. *et al.*. (2015) Multilocular Pancreatic Acinar Cystadenoma Containing Areas of Multifocal Branch-Duct Intraductal Papillary Mucinous Neoplasm <u>J. Pancreas 15 (5):</u> 475-80.

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee

12 months from date of despatch

Health And Safety
Information

Material Safety Datasheet documentation #10040 available at:
10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...) HRP
Rabbit Anti Mouse IgG (STAR12...) RPE

Rabbit Anti Mouse IgG (STAR8...) <u>DyLight®800</u>
Human Anti Mouse IgG3 (HCA039...) <u>FITC, HRP</u>

Goat Anti Mouse IgG (STAR76...) RPE

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Goat Anti Mouse IgG IgA IgM (STAR87...) Alk. Phos., HRP

Rabbit Anti Mouse IgG (STAR13...) HRP
Goat Anti Mouse IgG (STAR70...) FITC
Rabbit Anti Mouse IgG (STAR9...) FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M388976:210806'

Printed on 29 Aug 2021