

Datasheet: 2150-1470 BATCH NUMBER 157902

Description:	RABBIT ANTI MOUSE COLLAGEN IV
Specificity:	COLLAGEN IV
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 ml

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
		Yes	No	Not Determined	Suggested Dilution		
	Immunohistology - Frozen				1/500		
	Immunohistology - Paraffin	-			1/500		
	ELISA	•			1/2000		
	Immunoprecipitation						
	Western Blotting			•			
	Immunofluorescence	-			1/40		
	Where this product has r	ot been t	ested for	use in a particular tech	nnique this does not		
Target Species	a guide only. It is recomn system using appropriate Mouse			•	or use in their own		
Species Cross Reactivity	Reacts with: Orangutan, N.B. Antibody reactivity a reactivity is derived from personal communications further information.	and workin testing w	ng conditi ithin our la	aboratories, peer-revie	wed publications or		
Product Form	Purified Ig - liquid						
Preparation	Purified IgG prepared by	antigen c	olumn ch	romatography			
Buffer Solution	Phosphate buffered salin	е					

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃) Antibiotic antimycotic mixture 1%			
Immunogen	Collagen IV purified from mouse EHS tumor.			
External Database Links	UniProt:			
	P02463 Related reagents			
	P08122 Related reagents			
	Q9QZS0 Related reagents			
	Q9QZR9 Related reagents			
	Q80V57 Related reagents			
	Q6PFB1 Related reagents			
	Entrez Gene:			
	12826 Col4a1 Related reagents			
	12827 Col4a2 Related reagents			
	12828 Col4a3 Related reagents			
	12829 Col4a4 Related reagents			
RRID	AB_2082660			
	 IV is a 1682 amino acid ~160 kDa (predicted) matrx protein and major component of glomerular basement membranes. Multiple isoforms exist each capable of forming triple helical structures with two other chains to form the type IV collagen network. The collagen IV alpha chain can be cleaved between residues 1444-1445 to yield the c-terminal 225 amino acid, ~28 kDa arresten fragment, collagen α2(IV) yields a c-terminal canstatin fragment while Collagen α3(IV) yields a tumstatin fragment. Collagen IV bears a single collagen IV NC1 (C-terminal non-collagenous) domain (UniProt: Q9QZR9). Mutations in collagen IV genes have been implicated in inherited nephropathies and potentially in cystic kidney disease and intracranial aneurysms (Plaisier <i>et al.</i> 2007). 			
	Rabbit amti Mouse Collagen IV antibody has been successfully employed for the detection of collagen IV by immunofluorescence and immunohistochemistry in mice (<u>Tang <i>et al.</i></u> 2010), rats (<u>Shen <i>et al.</i> 2014</u>) and oragutan (<u>Bredies <i>et al.</i> 2013</u>).			
	The following cross reactivities have been observed:			
	Mouse type IV 100%			
	Mouse types I, II & III <0.1%			
	Human types IV & V <0.1%			
	Mouse fibronectin & laminin <0.1%			
References	1. Tang, Z. et al. (2010) Survival effect of PDGF-CC rescues neurons from apoptosis in			

both brain and retina by regulating GSK3beta phosphorylation. <u>J Exp Med. 207: 867-80.</u> 2. Fantin, A. *et al.* (2010) Tissue macrophages act as cellular chaperones for vascular anastomosis downstream of VEGF-mediated endothelial tip cell induction. <u>Blood.116:</u> 829-40.

3. Shen, W. *et al.* (2014) Systemic Administration of Erythropoietin Inhibits Retinopathy in RCS Rats. <u>PLoS One. 9: e104759.</u>

4. Bredies, K. *et al.* (2013) Computer-assisted counting of retinal cells by automatic segmentation after TV denoising. <u>BMC Ophthalmol. 13: 59.</u>

5. Rubin, A.N. *et al.* (2010) The germinal zones of the basal ganglia but not the septum generate GABAergic interneurons for the cortex. <u>J Neurosci. 30: 12050-62.</u>

6. Weinl, C. *et al.* (2014) Elk3 deficiency causes transient impairment in post-natal retinal vascular development and formation of tortuous arteries in adult murine retinae. <u>PLoS</u> <u>One. 9: e107048.</u>

7. Stenzel, D. *et al.* (2011) Integrin-dependent and -independent functions of astrocytic fibronectin in retinal angiogenesis. <u>Development. 138: 4451-63.</u>

8. Zuercher, J. *et al.* (2012) Norrin stimulates cell proliferation in the superficial retinal vascular plexus and is pivotal for the recruitment of mural cells. <u>Hum Mol Genet. 21:</u> 2619-30.

9. Arnold, T.D. *et al.* (2012) Defective retinal vascular endothelial cell development as a consequence of impaired integrin $\alpha V\beta 8$ -mediated activation of transforming growth factor- β . J Neurosci. 32: 1197-206.

10. Chen, M. *et al.* (2010) Immune activation in retinal aging: a gene expression study. Invest Ophthalmol Vis Sci. 51: 5888-96.

11. Kojima, T. *et al.* (2007) Proangiogenic role of ephrinB1/EphB1 in basic fibroblast growth factor-induced corneal angiogenesis. <u>Am J Pathol. 170: 764-73.</u>

12. Ma, S. *et al.* (2012) Ric-8a, a Guanine Nucleotide Exchange Factor for Heterotrimeric G Proteins, Regulates Bergmann Glia-Basement Membrane Adhesion during Cerebellar Foliation. <u>J Neurosci. 32: 14979-93.</u>

13. Dulauroy, S. *et al.* (2012) Lineage tracing and genetic ablation of ADAM12(+) perivascular cells identify a major source of profibrotic cells during acute tissue injury. <u>Nat</u> <u>Med. 18: 1262-70.</u>

14. Edgar, K. *et al.* (2012) eNOS Overexpression Exacerbates Vascular Closure in the Obliterative Phase of OIR and Increases Angiogenic Drive in the Subsequent Proliferative Stage. <u>Invest Ophthalmol Vis Sci. 53: 6833-50.</u>

15. Lange, C.A. *et al.* (2012) Von Hippel-Lindau protein in the RPE is essential for normal ocular growth and vascular development. <u>Development. 139: 2340-50.</u>

16. Li, W. and Mukouyama, Y.S. (2011) Whole-mount immunohistochemical analysis for embryonic limb skin vasculature: a model system to study vascular branching morphogenesis in embryo. J Vis Exp. 51: pii: 2620.

17. Lutter, S. *et al.* (2012) Smooth muscle-endothelial cell communication activates Reelin signaling and regulates lymphatic vessel formation. <u>J Cell Biol. 197: 837-49.</u>

18. McKenzie, J.A. *et al.* (2012) Apelin is required for non-neovascular remodeling in the retina. <u>Am J Pathol. 180: 399-409.</u>

19. Powner, M.B. *et al.* (2012) Visualization of gene expression in whole mouse retina by in situ hybridization. <u>Nat Protoc. 7: 1086-96.</u>

20. Schulz, C. *et al.* (2012) A lineage of myeloid cells independent of Myb and hematopoietic stem cells. <u>Science. 336: 86-90.</u>

21. Scott, A. *et al.* (2010) Astrocyte-derived vascular endothelial growth factor stabilizes vessels in the developing retinal vasculature. <u>PLoS One. 5: e11863.</u>

22. Takagi, N. *et al.* (2012) Mineralocorticoid Receptor Blocker Protects against Podocyte-Dependent Glomerulosclerosis. <u>Nephron Extra. 2: 17-26.</u>

23. Chen, M. *et al.* (2013) Age- and light-dependent development of localised retinal atrophy in CCL2(-/-)CX3CR1(GFP/GFP) mice. <u>PLoS One. 8: e61381.</u>

24. Luhmann UF. *et al.* (2005) Berger W. Role of the Norrie disease pseudoglioma gene in sprouting angiogenesis during development of the retinal vasculature. <u>Invest Ophthalmol</u> <u>Vis Sci. 46: 3372-82.</u>

25. Scott, A. *et al.* (2014) Quantification of vascular tortuosity as an early outcome measure in oxygen induced retinopathy (OIR) <u>Exp Eye Res. 120: 55-60.</u>

26. Yukiura H *et al.* (2015) Autotaxin overexpression causes embryonic lethality and vascular defects. <u>PLoS One. 10 (5): e0126734.</u>

27. Williams, J.A. *et al.* (2016) Regulation of C3 Activation by the Alternative Complement Pathway in the Mouse Retina. <u>PLoS One. 11 (8): e0161898.</u>

 28. Piñero G *et al.* (2016) Lithium Reversibly Inhibits Schwann Cell Proliferation and Differentiation Without Inducing Myelin Loss. <u>Mol Neurobiol. Dec 5. [Epub ahead of print]</u>
 29. Gurnik, S. *et al.* (2016) Angiopoietin-2-induced blood-brain barrier compromise and increased stroke size are rescued by VE-PTP-dependent restoration of Tie2 signaling. Acta Neuropathol. 131 (5): 753-73.

30. Misra, A. *et al.* (2016) Integrin β 3 inhibition is a therapeutic strategy for supravalvular aortic stenosis. <u>J Exp Med. 213 (3): 451-63.</u>

31. Wu, W.K. *et al.* (2015) IL-4 regulates specific Arg-1(+) macrophage sFIt-1-mediated inhibition of angiogenesis. <u>Am J Pathol. 185 (8): 2324-35.</u>

32. Toffoli, B. *et al.* (2017) Nephropathy in Pparg-null mice highlights PPARγ systemic activities in metabolism and in the immune system. <u>PLoS One. 12 (2): e0171474.</u>

33. Yanagida, K. *et al.* (2017) Size-selective opening of the blood-brain barrier by targeting endothelial sphingosine 1-phosphate receptor 1. <u>Proc Natl Acad Sci U S A. 114 (17)</u>: 4531-6.

34. Fernández-robredo, P. *et al.* (2017) Neuropilin 1 Involvement in Choroidal and Retinal Neovascularisation. <u>PLoS One. 12 (1): e0169865.</u>

35. Kim, B. *et al.* (2018) Endothelial pyruvate kinase M2 maintains vascular integrity. J <u>Clin Invest. 128 (10): 4543-56.</u>

36. Villard, O. *et al.* (2020) Characterization of immortalized human islet stromal cells reveals a MSC-like profile with pancreatic features. <u>Stem Cell Res Ther. 11 (1): 158.</u>

37. Chu, C.J. *et al.* (2013) Assessment and *in vivo* scoring of murine experimental autoimmune uveoretinitis using optical coherence tomography. <u>PLoS One. 8 (5): e63002.</u>
38. Krivanek, J. *et al.* (2020) Dental cell type atlas reveals stem and differentiated cell types in mouse and human teeth. <u>Nat Commun. 11 (1): 4816.</u>

39. Elabi, O. *et al.* (2021) Human α -synuclein overexpression in a mouse model of Parkinson's disease leads to vascular pathology, blood brain barrier leakage and pericyte activation. <u>Sci Rep. 11 (1): 1120.</u>

Storage

Store at +4°C or at -20°C if preferred.

Storage in frost-free freezers is not recommended.

This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend

	microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #20470 available at: https://www.bio-rad-antibodies.com/SDS/2150-1470 20470
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (Fc) (STAR121...)Biotin, FITC, HRPSheep Anti Rabbit IgG (STAR35...)RPEGoat Anti Rabbit IgG (H/L) (STAR124...)HRP

Recommended Useful Reagents

ANTIGEN RETRIEVAL BUFFER, pH8.0 (BUF025A)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	id.com	Email: antibody_sales_uk@bio-ra	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M372167:200630'

Printed on 16 Apr 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint