

# Datasheet: 2485-4906

Description:	SHEEP ANTI YEAST CYTOSINE DEAMINASE
Specificity:	CYTOSINE DEAMINASE
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 ml

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .							
		Yes	No	Not Determined	Suggested Dilution			
	ELISA	•			1/500 - 1/2500			
	Western Blotting	•			1/100 - 1/500			
	Where this product has not been tested for use in a particular technique this does not							
	necessarily exclude its u a guide only. It is recomr system using appropriate	mended th	at the use	er titrates the product f	• •			
Target Species	Yeast							
Product Form	Purified Ig - liquid							
Preparation	Purified Ig prepared by affinity chromatography on Protein G							
Antiserum Preparatio	n Antisera to cytosine dea highly purified antigen.	minase we	ere raised	by repeated immunisa	ations sheep of with			
Buffer Solution	Phosphate buffered salir	ne						
Preservative Stabilisers	0.09% Sodium Azide (Na	aN <sub>3</sub> )						
Approx. Protein Concentrations	lg concentration 5mg/ml							
Immunogen	Purified yeast cytosine d	eaminase	•					

External Database Links	UniProt: <u>Q12178</u> Related reagents							
RRID	AB_618795							
Specificity	Sheep anti Yeast Cytosine Deaminase antibody recognizes cytosine deaminase, also known as cytosine aminohydrolase. Cytosine deaminase is a 158 amino acid ~18 kDa enzyme involved in the conversion of cytosine to uracil. Sheep anti Yeast Cytosine Deaminase antibody recognizes the native antigen in ELISA and has been successfully used for detection of the enzyme by western blotting (Fuerer and Iggo 2004).							
Western Blotting	In Western blot two bands were visualised by ECL at 13kD and 17kD.							
References	<ol> <li>Pan, X. <i>et al.</i> (2003) Adenovirus-mediated gene transfer in the treatment of pancreatic cancer. Pancreas. 26: 274-8.</li> <li>Fuerer, C. <i>et al.</i> (2004) 5-Fluorocytosine increases the toxicity of Wnt-targeting replicating adenoviruses that express cytosine deaminase as a late gene. Gene Ther. 11: 142-51</li> <li>Conrad, C. <i>et al.</i> (2005) Δ24-hyCD adenovirus suppresses glioma growth <i>in vivo</i> by combining oncolysis and chemosensitization. Cancer Gene Ther. 12: 284-94.</li> <li>Michnick, S.W. <i>et al.</i> (2010) A toolkit of protein-fragment complementation assays for studying and dissecting large-scale and dynamic protein-protein interactions in living cells. Methods Enzymol. 470: 335-68.</li> <li>Hammer, K. <i>et al.</i> (2015) Engineered adenoviruses combine enhanced oncolysis with improved virus production by mesenchymal stromal carrier cells. Int J Cancer. 137 (4): 978-90.</li> <li>Komissarov, A. <i>et al.</i> (2017) Cytotoxic effect of co-expression of human hepatitis A virus 3C protease and bifunctional suicide protein FCU1 genes in a bicistronic vector. Mol Biol Rep. 44 (4): 323-32.</li> <li>Hanauer, J.R.H. <i>et al.</i> (2019) High-affinity DARPin allows targeting of measles virus to glioblastoma multiforme in combination with protease targeting without loss of potency. Molecular Therapy - Oncolytics. 14. Oct [Epub ahead of print].</li> </ol>							
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.							
Guarantee	12 months from date of despatch							
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/2485-4906 10040							
Regulatory	For research purposes only							

### **Related Products**

#### **Recommended Secondary Antibodies**

Rabbit Anti Sheep IgG (H/L) (5184-2304...) Biotin

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	ad.com	Email: antibody_sales_uk@bio-ra	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M429904:240501'

#### Printed on 23 May 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint