Datasheet: 2722-5204

Description:	MOUSE ANTI DESMOPLAKIN 1/2
Specificity:	DESMOPLAKIN 1/2
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	DP-2.15
Isotype:	IgG1
Quantity:	50 µg

Product Details

RRID AB_619950

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	▪			
Western Blotting	▪			
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

Target Species Human

Species Cross Reactivity Reacts with: Bovine, Chicken, Mouse, Rat
N.B. Antibody reactivity and working conditions may vary between species.

Product Form Purified IgG - liquid

Preparation Purified IgG prepared by affinity chromatography on Protein A

Buffer Solution Phosphate buffered saline

Preservative Stabilisers 0.09% Sodium Azide (NaN_3)

Approx. Protein Concentrations IgG concentration 1mg/ml

Immunogen Native bovine desmoplakin 1 and 2 purified from bovine snout desmosomes.

External Database

UniProt:

Links	P15924 Related reagents **Entrez Gene:** 1832 DSP Related reagents
Fusion Partners	Spleen cells from immunized Balb/c mice were fused with cells from the NS0 mouse myeloma cell line.
Specificity	**Mouse anti Desmoplakin 1/2 antibody, clone DP-2.15** recognizes both desmoplakin 1 and 2 from stratified epithelia, simple epithelia including glands, urothelium, thymic reticular epithelium, hepatocytes, intercalated disks of myocardium and arachnoid cells of meninges. Mouse anti Desmoplakin 1/2 antibody, clone DP-2.15 can be used for the detection of primary and metastatic carcinomas. Mouse anti Desmoplakin 1/2 antibody, clone DP-2.15 recognizes the N terminus of the rod domain of desmoplakin (Bornslaeger *et al.* 1996).
References	1. Cowin, P. *et al.* (1985) The complement of desmosomal plaque proteins in different cell types. J Cell Biol. 101 (4): 1442-54. 2. Jheon, A.H. *et al.* (2011) PERP regulates enamel formation via effects on cell-cell adhesion and gene expression. J Cell Sci. 124: 745-54. 3. Bornslaeger, E.A. *et al.* (1996) Breaking the connection: displacement of the desmosomal plaque protein desmoplakin from cell-cell interfaces disrupts anchorage of intermediate filament bundles and alters intercellular junction assembly. J. Cell. Biol. 134: 985-1001. 4. Park, D.S. *et al.* (2015) Genetically engineered SCN5A mutant pig hearts exhibit conduction defects and arrhythmias. J Clin Invest. 125 (1): 403-12. 5. DeQuach JA *et al.* (2010) Simple and high yielding method for preparing tissue specific extracellular matrix coatings for cell culture. PLoS One. 5 (9): e13039. 6. Basheer, W.A. *et al.* (2015) Cardiomyocyte-specific overexpression of the ubiquitin ligase Wwp1 contributes to reduction in Connexin 43 and arrhythmogenesis. J Mol Cell Cardiol. 88: 1-13. 7. Yung, H.W. *et al.* (2016) Placental endoplasmic reticulum stress in gestational diabetes: the potential for therapeutic intervention with chemical chaperones and antioxidants. Diabetologia. 59 (10): 2240-50. 8. Chkourko, H.S. *et al.* (2012) Remodeling of mechanical junctions and of microtubule-associated proteins accompany cardiac connexin43 lateralization. Heart Rhythm. 9 (7): 1133-1140.e6. 9. Vila, J. *et al.* (2017) Structural and molecular pathology of the atrium in boxer arrhythmogenic right ventricular cardiomyopathy. J Vet Cardiol. 19 (1): 57-67. 10. Yung, H.W. *et al.* (2016) Placental endoplasmic reticulum stress in gestational diabetes: the potential for therapeutic intervention with chemical chaperones and antioxidants. Diabetologia. 59 (10): 2240-50.
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG IgA IgM (STAR87...)	Alk. Phos., HRP
Goat Anti Mouse IgG (STAR77...)	HRP
Rabbit Anti Mouse IgG (STAR12...)	RPE
Rabbit Anti Mouse IgG (STAR8...)	DyLight®800
Rabbit Anti Mouse IgG (STAR13...)	HRP
Goat Anti Mouse IgG (STAR76...)	RPE
Goat Anti Mouse IgG (STAR70...)	FITC
Goat Anti Mouse IgG (Fc) (STAR120...)	FITC, HRP
Rabbit Anti Mouse IgG (STAR9...)	FITC
Goat Anti Mouse IgG (H/L) (STAR117...)	Alk. Phos., DyLight®488, DyLight®549, DyLight®649, DyLight®680, DyLight®800, FITC, HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South America
Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide
Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe
Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

'M336798:181214'

Printed on 15 Mar 2019