

## Datasheet: 3710-0956

<b>Description:</b>	GOAT ANTI DIPHTHERIA TOXIN
<b>Specificity:</b>	DIPHTHERIA TOXIN
<b>Format:</b>	Purified
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	1 ml

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
ELISA	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

<b>Target Species</b>	Bacterial
<b>Product Form</b>	Purified IgG - liquid
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.1% Sodium Azide (NaN <sub>3</sub> )
<b>Approx. Protein Concentrations</b>	4 mg/ml
<b>Immunogen</b>	Purified diphtheria toxoid
<b>External Database Links</b>	<b>UniProt:</b> <a href="#">Q6NK15</a> <a href="#">Related reagents</a>
<b>RRID</b>	AB_620528

**Specificity**

**Goat anti diphtheria toxin antibody** reacts with the diphtheria toxin/toxoid.

Diphtheria toxin is an exotoxin secreted by various strains of *Corynebacterium diphtheriae* that is the cause of diphtheria. The active toxin is a single chain of 535 amino acids composed of 3 distinct structural/functional domains; and N-Terminal ADP-ribosyltransferase (catalytic domain); a region which facilitates the delivery of the catalytic domain across the cell membrane (transmembrane domain) and the eukaryotic cell receptor binding domain.

Goat anti diphtheria toxin antibody also recognizes CRM<sub>197</sub> ([Conde et al.2021](#)), a non-toxic mutant form of diphtheria toxin with an amino acid substitution at glycine-52 to glutamic acid ([Giannini et al. 1984](#)).

**References**

1. Conde, E. *et al.* (2021) Dual vaccination against IL-4 and IL-13 protects against chronic allergic asthma in mice. [Nat Commun. 12 \(1\): 2574.](#)

**Storage**

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

**Guarantee**

12 months from date of despatch

**Health And Safety Information**

Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

**Regulatory**

For research purposes only

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) [FITC](#), [HRP](#)

**North & South America** Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M382493:210513'

Printed on 29 Aug 2021