

# Datasheet: 4050-0096 BATCH NUMBER 164762

Description:	MOUSE ANTI HUMAN ELASTASE		
Specificity:	city: ELASTASE		
Other names:	ELA2		
Format:	Purified		
Product Type:	Monoclonal Antibody		
Clone:	ELA10-101.5 (39A)		
Isotype:	lgG1		
Quantity:	0.2 mg		

## **Product Details**

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				
Immunohistology - Frozen			•	
Immunohistology - Paraffin				
ELISA				
Western Blotting			•	
Immunofluorescence				
Functional Assays				

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

Target Species	Human	
Product Form	Purified IgG - liquid	
Preparation	Purified IgG prepared by affinity chromatography on Protein G	from ascites
Buffer Solution	Phosphate buffered saline	
Preservative	<0.1% Sodium Azide (NaN <sub>3</sub> )	

### Stabilisers **Approx. Protein** 0.5 mg/ml Concentrations Immunogen Elastase from human neutrophils. **External Database UniProt:** Links P08246 Related reagents Entrez Gene: 1991 ELANE Related reagents Synonyms ELA2 RRID AB 617258 Specificity Mouse anti Human elastase antibody, clone 39A recognizes human neutrophil elastase, also known as elastase-2, Bone marrow serine protease, PMN elastase or Medullasin. Elastase is a 238 amino acid ~27 kDa enzyme with an addiional 29 amino acid signal and pro-peptide region. Mutations of the ELA2 gene can lead to a number of hematopoietic disorders including cyclic hematopoiesis (CH) characterized by a 21 day cycle of bone marrow derived blood cell production with neutrophil numbers ranging between normal levels and almost zero (Horwitz et al. 1999) and severe congential autosomal neutropenia (SCN1) demonstating developmental arrest of granulopoiesis at the promyelocyte stage (Bellanné-Chantelot et al. 2004) with both conditions leading to a predisposition to bacterial infections .. References 1. Buján, J. et al. (2003) Expression of elastic components in healthy and varicose veins. World J Surg. 27 (8): 901-5. 2. Durant, S. et al. (2004) Apoptosis-induced proteinase 3 membrane expression is independent from degranulation. J Leukoc Biol. 75 (1): 87-98. 3. Korkmaz, B. et al. (2005) Inhibition of neutrophil elastase by alpha1-protease inhibitor at the surface of human polymorphonuclear neutrophils. J Immunol. 175 (5): 3329-38. 4. Vong, L. et al. (2007) Annexin 1 cleavage in activated neutrophils: a pivotal role for proteinase 3. J Biol Chem. 282 (41): 29998-30004. 5. Davies, P.L. et al. (2008) Monoclonal anti-neutrophil elastase antibody characterisation: ability to block function, detect free versus serpin-complexed enzyme and stain intracellular granules. J Immunol Methods. 336 (2): 175-82. 6. Muthana, M. et al. (2015) C5orf30 is a negative regulator of tissue damage in rheumatoid arthritis. Proc Natl Acad Sci U S A. 112 (37): 11618-23. Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch				
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/4050-0096 10040				
Regulatory	For research purposes only				

# Related Products

### **Recommended Secondary Antibodies**

Goat Anti Mouse IgG (STAR77)	HRP			
Rabbit Anti Mouse IgG (STAR12)	RPE			
Goat Anti Mouse IgG IgA IgM (STAR87) <u>Alk. Phos.</u> , <u>HRP</u>				
Goat Anti Mouse IgG (STAR76)	RPE			
Rabbit Anti Mouse IgG (STAR13)	HRP			
Goat Anti Mouse IgG (STAR70)	FITC			
Goat Anti Mouse IgG (H/L) (STAR117)	Alk. Phos., DyLight®488, DyLight®550,			
	<u>DyLight®650</u> , <u>DyLight®680</u> , <u>DyLight®800</u> ,			
	<u>FITC</u> , <u>HRP</u>			
Rabbit Anti Mouse IgG (STAR9)	FITC			
Goat Anti Mouse IgG (Fc) (STAR120)	<u>FITC</u> , <u>HRP</u>			
<b>Recommended Negative Controls</b>				

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	ad.com	Email: antibody_sales_uk@bio-i	ad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M405735:220916'

#### Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint