

Datasheet: 4510-7804

**BATCH NUMBER 170092**

<b>Description:</b>	RABBIT ANTI FITC
<b>Specificity:</b>	FITC
<b>Other names:</b>	FLUORESCEIN ISOTHIOCYANATE
<b>Format:</b>	Purified
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
ELISA	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

<b>Target Species</b>	Chemical
<b>Product Form</b>	Purified Ig - liquid
<b>Preparation</b>	Purified IgG prepared by immunoaffinity chromatography
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.1% Sodium Azide (NaN <sub>3</sub> )
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	FITC Isomer 1.

---

**Specificity** **Rabbit anti FITC antibody** recognizes fluorescein and does not react with immunoglobulins from any species.

Fluorescein isothiocyanate (FITC) is widely used to attach a fluorescent label to proteins via the amine group. The isothiocyanate group reacts with amino terminal and primary amines in proteins.

---

**References**

1. van der Loos, C. (2008) Multiple immunoenzyme staining: methods and visualizations for the observation with spectral imaging. [J. Histochem Cytochem. 56: 313-328.](#)
2. Bdeir, K. *et al.* (2010) Neutrophil alpha-defensins cause lung injury by disrupting the capillary-epithelial barrier. [Am J Respir Crit Care Med. 181: 935-46.](#)
3. Hsiao, J.R. *et al.* (2010) Cyclic alphavbeta6-targeting peptide selected from biopanning with clinical potential for head and neck squamous cell carcinoma. [Head Neck. 32 \(2\): 160-72.](#)
4. Türeci, ö *et al.* (2011) Claudin-18 gene structure, regulation, and expression is evolutionary conserved in mammals. [Gene. 481 \(2\): 83-92.](#)
5. Martínez-Sernández, V. *et al.* (2011) Development and evaluation of a new lateral flow immunoassay for serodiagnosis of human fasciolosis. [PLoS Negl Trop Dis. 5 \(11\): e1376.](#)
6. Martínez-Sernández, V. *et al.* (2014) The MF6p/FhHDM-1 major antigen secreted by the trematode parasite *Fasciola hepatica* is a heme-binding protein. [J Biol Chem. 289 \(3\): 1441-56.](#)
7. Dileepan, T. *et al.* (2015) *In Situ* Peptide-MHC-II Tetramer Staining of Antigen-Specific CD4+ T Cells in Tissues. [PLoS One. 10 \(6\): e0128862.](#)
8. Arni, S. *et al.* (2021) Subnormothermic *ex vivo* lung perfusion attenuates ischemia reperfusion injury from donation after circulatory death donors. [PLoS One. 16 \(8\): e0255155.](#)

---

**Storage**

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety Information**

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/4510-7804>  
10040

---

**Regulatory**

For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...) [FITC](#)

Goat Anti Rabbit IgG (H/L) (STAR124...) [HRP](#)  
Sheep Anti Rabbit IgG (STAR35...) [RPE](#)  
Goat Anti Rabbit IgG (Fc) (STAR121...) [Biotin](#), [FITC](#), [HRP](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)

'M381432:210512'

Printed on 17 Mar 2025

---

© 2025 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)