

Datasheet: 4745-1051

Description:	SHEEP ANTI GREEN FLUORESCENT PROTEIN
Specificity:	GREEN FLUORESCENT PROTEIN
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			
Western Blotting			▪	
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G.
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 5.0 mg/ml
Immunogen	Green fluorescent protein from <i>Aequorea victoria</i> .
External Database Links	UniProt:

RRID AB_619712

Specificity **Sheep anti Green Fluorescent Protein antibody** recognizes green fluorescent protein (GFP), a ~27 kDa protein derived from the jellyfish *Aequorea victoria*. GFP fluoresces green (509nm) when excited by blue light (395nm) and is commonly used as a marker of gene expression.

- References**
1. Collins, R.T. *et al.* (2010) MAZe: a tool for mosaic analysis of gene function in zebrafish. [Nat Methods. 7: 219-23.](#)
 2. Wu, L. *et al.* (2011) Properties of a distinct subpopulation of GABAergic commissural interneurons that are part of the locomotor circuitry in the neonatal spinal cord. [J Neurosci. 31 \(13\): 4821-33.](#)
 3. Knipe, L. *et al.* (2010) A revised model for the secretion of tPA and cytokines from cultured endothelial cells. [Blood. 116: 2183-91.](#)
 4. Shneider, N.A. *et al.* (2009) Gamma motor neurons express distinct genetic markers at birth and require muscle spindle-derived GDNF for postnatal survival. [Neural Dev. 4: 42.](#)
 5. Soza-Ried, C. *et al.* (2008) Maintenance of thymic epithelial phenotype requires extrinsic signals in mouse and zebrafish. [J Immunol. 181: 5272-7.](#)
 6. Lopez, K.A. *et al.* (2011) Convection-enhanced delivery of topotecan into a PDGF-driven model of glioblastoma prolongs survival and ablates both tumor-initiating cells and recruited glial progenitors. [Cancer Res. 71: 3963-71.](#)
 7. League, G.P. and Nam, S.C. (2011) Role of kinesin heavy chain in Crumbs localization along the rhabdome elongation in *Drosophila* photoreceptor. [PLoS One. 6:e21218.](#)
 8. Siembab, V.C. *et al.* (2010) Target selection of proprioceptive and motor axon synapses on neonatal V1-derived Ia inhibitory interneurons and Renshaw cells. [J Comp Neurol. 518: 4675-701.](#)
 9. Srinivasan, S. *et al.* (2012) The receptor tyrosine phosphatase Lar regulates adhesion between *Drosophila* male germline stem cells and the niche. [Development. 139: 1381-90.](#)
 10. Haberlandt, C. *et al.* (2011) Gray matter NG2 cells display multiple Ca²⁺-signaling pathways and highly motile processes. [PLoS One. 6: e17575.](#)
 11. Cheung, L.S. *et al.* (2013) Dynamic model for the coordination of two enhancers of broad by EGFR signaling. [Proc Natl Acad Sci U S A. 110: 17939-44.](#)
 12. Li, X. *et al.* (2013) Temporal patterning of *Drosophila* medulla neuroblasts controls neural fates. [Nature. 498: 456-62.](#)
 13. Behnia, R. *et al.* (2014) Processing properties of ON and OFF pathways for *Drosophila* motion detection. [Nature. 512: 427-30.](#)
 14. de Nooij, J.C. *et al.* (2015) The PDZ-domain protein Whirlin facilitates mechanosensory signaling in mammalian proprioceptors. [J Neurosci. 35 \(7\): 3073-84.](#)
 15. Scotti, M. *et al.* (2015) A Hoxa13:Cre mouse strain for conditional gene manipulation in developing limb, hindgut, and urogenital system. [Genesis. 53 \(6\): 366-76.](#)
 16. Sun, G.J. *et al.* (2015) Latent tri-lineage potential of adult hippocampal neural stem cells revealed by Nf1 inactivation. [Nat Neurosci. 18 \(12\): 1722-4.](#)
 17. Schlegel, P. *et al.* (2016) Synaptic transmission parallels neuromodulation in a central food-intake circuit. [eLife 2016;10.7554/eLife.16799](#)
 18. Crouch, E.E. *et al.* (2015) Regional and stage-specific effects of prospectively purified

- vascular cells on the adult V-SVZ neural stem cell lineage. [J Neurosci. 35 \(11\): 4528-39.](#)
19. Gushchina, S. *et al.* (2018) Increased expression of colony-stimulating factor-1 in mouse spinal cord with experimental autoimmune encephalomyelitis correlates with microglial activation and neuronal loss. [Glia. 66 \(10\): 2108-25.](#)
20. Won, J.H. *et al.* (2019) ADAMTS Sol narae cleaves extracellular Wingless to generate a novel active form that regulates cell proliferation in *Drosophila*. [Cell Death Dis. 10 \(8\): 564.](#)
21. Del Valle Rodríguez, A. *et al.* (2020) A network approach to analyze neuronal lineage and layer innervation in the *Drosophila* optic lobes. [PLoS One. 15 \(2\): e0227897.](#)
22. Choquet, C. *et al.* (2020) Nkx2-5 defines distinct scaffold and recruitment phases during formation of the murine cardiac Purkinje fiber network. [Nat Commun. 11 \(1\): 5300.](#)
23. Lee, S.R. *et al.* (2020) Regulation of epithelial integrity and organ growth by Tctp and Coracle in *Drosophila*. [PLoS Genet. 16 \(6\): e1008885.](#)
24. Oliver, K.M. *et al.* (2021) Molecular correlates of muscle spindle and Golgi tendon organ afferents. [Nat Commun. 12 \(1\): 1451.](#)

Further Reading 1. Adams, K.L. *et al.* (2015) Foxp1-mediated programming of limb-innervating motor neurons from mouse and human embryonic stem cells. [Nat Commun. 6: 6778.](#)

Storage Store at +4°C or at -20°C if preferred.
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Sheep IgG (H/L) (5184-2304...) [Biotin](#)

Donkey Anti Sheep IgG (STAR88...) [DyLight®488](#), [HRP](#)

Recommended Useful Reagents

[RABBIT ANTI BLUE FLUORESCENT PROTEIN \(AHP2985\)](#)

[RABBIT ANTI CYAN FLUORESCENT PROTEIN \(AHP2986\)](#)

[RABBIT ANTI RED FLUORESCENT PROTEIN \(AHP2987\)](#)

[MOUSE ANTI mCHERRY \(MCA6020\)](#)

[RABBIT ANTI mCHERRY \(AHP2326\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751
Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739
Email: antibody_sales_uk@bio-rad.com

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50
Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M363150:200528'

Printed on 23 Mar 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)