

Datasheet: 7460-3104 **BATCH NUMBER 165280**

Description:	MOUSE ANTI HUMAN CD140b		
Specificity:	CD140b		
Other names:	PDGF RECEPTOR BETA		
Format:	Purified		
Product Type:	Monoclonal Antibody		
Clone:	PR7212		
Isotype:	lgG1		
Quantity:	0.1 mg		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			25 🍫 g/ml
Immunohistology - Frozen	•			
Immunohistology - Paraffin	•			25 🍫g/ml
Immunoprecipitation	•			
Western Blotting	•			
Immunofluorescence	•			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

Target Species	Human
Species Cross	Reacts with: Baboon, Monkey
Reactivity	Does not react with:Mouse, Rat
	N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG - liquid

Preparation	Purified IgG prepared by affinity chromatography on Protein G from ascites
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	1.0 mg/ml
Immunogen	Human skin fibroblast cell membrane extracts.
External Database Links	UniProt: P09619 Related reagents Entrez Gene: 5159 PDGFRB Related reagents
RRID	AB_2162788
Specificity	Mouse anti Human CD140b antibody, clone PR7212 recognizes the extracellular domain of the human PDGF receptor beta subunit., confirmed using several different cell lines and by its ability to immunoprecipitate PDGF receptor beta complexed with ¹²⁵ I-PDGF-BB (Hart et al. 1987). Mouse anti Human CD140b antibody, clone PR7212 does not recognize PDGF receptor alpha. PDGF receptor beta is a member of the class III receptor tyrosine kinase family that also includes M-CSF receptor, SCF receptor and FIt-3. Binding of PDGF-BB induces receptor homodimerization or heterodimerization with PDGF receptor alpha.
Histology Positive Control Tissue	Human breast cancer tissue
References	 Hart, C.E. et al. (1988) Two classes of PDGF receptor recognize different isoforms of PDGF. Science. 240: 1529-31. Seifert, R.A. et al. (1989) Two different subunits associate to create isoform-specific platelet-derived growth factor receptors. J Biol Chem. 264: 8771-8. Franklin, W.A. et al. (1990) In situ distribution of the beta-subunit of platelet-derived growth factor receptor in nonneoplastic tissue and in soft tissue tumors. Cancer Res. 50: 6344-8. Krane, J.F. et al. (1991) Increased dermal expression of platelet-derived growth factor receptors in growth-activated skin wounds and psoriasis. J Invest Dermatol. 96: 983-6. Palman, C. et al. (1992) Platelet-derived growth factor receptor (beta-subunit) immunoreactivity in soft tissue tumors. Lab Invest. 66: 108-15. Wang, J. et al. (1994) Cell proliferation in human soft tissue tumors correlates with platelet-derived growth factor B chain expression: an immunohistochemical and in situ hybridization study. Cancer Res. 54: 560-4. Ascoli, V. et al. (1995) Platelet-derived growth factor receptor immunoreactivity in

mesothelioma and nonneoplastic mesothelial cells in serous effusions. <u>Acta Cytol. 39:</u> 613-22.

8. Gilbertson, D.G. *et al.* (2001) Platelet-derived growth factor C (PDGF-C), a novel growth factor that binds to PDGF alpha and beta receptor. <u>J Biol Chem. 276: 27406-14.</u>

9. Meng, J. *et al.* (2011) Contribution of human muscle-derived cells to skeletal muscle regeneration in dystrophic host mice. PLoS One. 6: e17454.

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/7460-3104 10040
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (STAR77...) <u>HRP</u>

Rabbit Anti Mouse IgG (STAR12...) RPE

Goat Anti Mouse IgG IgA IgM (STAR87...) Alk. Phos., HRP

Goat Anti Mouse IgG (STAR76...) RPE

Goat Anti Mouse IgG (Fc) (STAR120...) FITC, HRP

Rabbit Anti Mouse IgG (STAR13...) HRP
Rabbit Anti Mouse IgG (STAR9...) FITC
Goat Anti Mouse IgG (STAR70...) FITC

Goat Anti Mouse IgG (H/L) (STAR117...) Alk. Phos., DyLight®488, DyLight®550,

DyLight®650, DyLight®680, DyLight®800,

FITC, HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South Tel: +1 800 265 7376 Worldwide Tel: +44 (0)1865 852 700

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

Europe

Tel: +49 (0) 89 8090 95 21

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M389059:210806'

Printed on 18 Jan 2024