

Datasheet: 7950-0004 BATCH NUMBER 167550

Description:	ption: GOAT ANTI RESPIRATORY SYNCYTIAL VIRUS			
Specificity:	RESPIRATORY SYNCYTIAL VIRUS			
Other names:	RSV			
Format:	Purified			
Product Type:	Polyclonal Antibody			
Isotype:	Polyclonal IgG			
Quantity:	1 ml			

Product Details

 Applications
 This product has been reported to work in the following applications. This information is
derived from testing within our laboratories, peer-reviewed publications or personal
communications from the originators. Please refer to references indicated for further
information. For general protocol recommendations, please visit www.bio-
rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			•	
Immunohistology - Frozen				
Immunohistology - Paraffin			•	
ELISA				
Immunofluorescence	•			
Functional Assays (1)				

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using the appropriate negative/positive controls.

(1)This product contains sodium azide, removal by dialysis is recommended prior to use in functional assays.

Target Species	Viral
Product Form	Purified IgG - liquid
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.1% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 5.0 mg/ml

Immunogen	Human RSV isolate.
RRID	AB_620536
Specificity	Goat anti respiratory syncitial virus polyclonal antibody recognizes respiratory syncytial virus (RSV) a negative-sense, single-stranded RNA virus and member of the <i>Paramyxoviridae</i> family. RSV causes respiratory tract infections in patients of all ages, but particularly affects infants and the immunosuppressed.
	RSV encodes three envelope glycoproteins, a small hydrophobic (SH) protein of unknown function, a glycoprotein (G) known as the attachment protein, and a fusion (F) protein. The F protein directs fusion of viral and cellular membranes, resulting in viral penetration, and can lead to the formation of syncytia.
	The F protein is thought to be the principal antigen responsible for inducing an immune response.
	Goat anti respiratory syncitial virus does not react with Parainfluenza 1-3, Influenza A and B, Adenovirus or uninfected HEp-2 or WI-38 cells. Goat anti respiratory syncitial virus polyclonal antibody is neutralizing and reacts well with bovine isolates.
References	 Culley, F.J. <i>et al.</i> (2006) Role of CCL5 (RANTES) in viral lung disease. J Virol. 80: 8151-7. Numata, M. <i>et al.</i> (2010) Pulmonary surfactant phosphatidylglycerol inhibits respiratory syncytial virus-induced inflammation and infection. Proc Natl Acad Sci U S A. 107: 320-5. Roux, X. <i>et al.</i> (2008) Sub-nucleocapsid nanoparticles: a nasal vaccine against respiratory syncytial virus. PLoS One. 3: e1766. Olszewska, W. <i>et al.</i> (2011) Antiviral and lung protective activity of a novel RSV fusion inhibitor in a mouse model. Eur Respir J. 38: 401-8. Fonceca AM <i>et al.</i> (2012) Primary airway epithelial cultures from children are highly permissive to respiratory syncytial virus infection. Thorax. 67 (1): 42-8. Ryzhakov, G. <i>et al.</i> (2011) IL-17 Boosts Proinflammatory Outcome of Antiviral Response in Human Cells. J Immunol. 187: 5357-62. Fricke J <i>et al.</i> (2013) p38 and OGT sequestration into viral inclusion bodies in cells infected with human respiratory syncytial virus suppresses MK2 activities and stress granule assembly. J Virol. 87 (3): 1333-47. Kipper, S. <i>et al.</i> (2015) New host factors important for respiratory syncytial virus (RSV) replication revealed by a novel microfluidics screen for interactors of matrix (M) protein. Mol Cell Proteomics. 14 (3): 532-43. Russell, R.F. <i>et al.</i> (2015) Partial Attenuation of Respiratory Syncytial Virus with a Deletion of a Small Hydrophobic Gene Is Associated with Elevated Interleukin-1β Responses. J Virol. 89 (17): 8974-81. Currie, S.M. <i>et al.</i> (2016) Cathelicidins Have Direct Antiviral Activity against Respiratory Syncytial Virus <i>In Vitro</i> and Protective Function <i>In Vivo</i> in Mice and Humans. J Immunol. 196 (6): 2699-710. Kinnear, E. <i>et al.</i> (2018) Airway T cells protect against RSV infection in the absence of antibody. Mucosal Immunol. 11 (1): 249-56.

	 Bajimaya, S. <i>et al.</i> (2017) Cholesterol is required for stability and infectivity of influenza A and respiratory syncytial viruses. <u>Virology. 510: 234-41.</u> Choi, E.J. <i>et al.</i> (2018) Exchange Proteins Directly Activated by cAMP and Their Roles in Respiratory Syncytial Virus Infection. <u>J Virol. 92 (22): e01200-18.</u> 		
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.		
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.		
Guarantee	12 months from date of despatch		
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/7950-0004 10040		
Regulatory	For research purposes only		

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) FITC, HRP

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50		
Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com			
To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M420461:230706'							

Printed on 10 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint