

Datasheet: 8450-0505

BATCH NUMBER 159319

Description:	RAT ANTI SUBSTANCE P
Specificity:	SUBSTANCE P
Format:	S/N
Product Type:	Monoclonal Antibody
Clone:	NC1/34
Isotype:	IgG2a
Quantity:	0.1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	▪			1/100 - 1/200
Immunohistology - Paraffin (1)	▪			1/100 - 1/200
Immunofluorescence	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

(1)This product does not require antigen retrieval prior to staining of formalin fixed, paraffin embedded sections.

Species Cross Reactivity

Reacts with: Pig, Pigeon, Bichir, Rabbit, Human, Bovine, Guinea Pig, Calanus sp., Cancer sp., Panilurus sp., Homarus sp., Mudpuppy, Lamprey, Crayfish, Zebrafish, Bullfrog, Cockroach, Sheep, Cowbird, Rat, Spadefoot toad, Chicken, Chinchilla, Camel, Beaver, Chinchilla

Based on sequence similarity, is expected to react with:Mammals, Crustacea, Amphibia
N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form	Tissue Culture Supernatant - liquid
---------------------	-------------------------------------

Preservative Stabilisers	0.05% Thiomersal
Immunogen	Substance P conjugated to Bovine Serum Albumin.
External Database Links	<p>UniProt: P20366 Related reagents</p> <p>Entrez Gene: 6863 TAC1 Related reagents</p>
Synonyms	NKA, NKNA, TAC2
RRID	AB_2200292
Fusion Partners	Spleen cells from immunised Wistar rat were fused with cells of the NS1/1-Ag 4-1 myeloma cell line.
Specificity	<p>Rat anti substance P antibody, clone NC1/34 recognizes the COOH terminal region of substance P, an eleven amino acid polypeptide neurotransmitter derived by proteolytic cleavage of protachykinin-1. Substance P binds to and activates the neurokinin-1 receptor. Substance P is a highly conserved neurotransmitter expressed throughout the animal kingdom. It is highly expressed in the spinal cord dorsal horn and dorsal root ganglia, but not in the ventral horn (Ribeiro-da-Silva and Hökfelt 2000). It is also expressed in non-neuronal cells including microglia and other immune cells.</p> <p>Substance P has a role in several physiologic activities such as pain transmission, the vomiting reflex, salivary secretion and smooth muscle contraction (Hökfelt 2004).</p> <p>Rat anti substance P antibody, clone NC1/34 demonstrates ~5% reactivity with eledoisin a neurotransmitter identified in the octopus Genus <i>Eledone</i>. It does not react with Leu or Met-enkephalin, somatostatin or beta-endorphin in tested applications.</p>
References	<ol style="list-style-type: none"> Kozsurek, M. <i>et al.</i> (2009) Nonselective innervation of lamina I projection neurons by cocaine- and amphetamine-regulated transcript peptide (CART)-immunoreactive fibres in the rat spinal dorsal horn. Eur J Neurosci. 29: 2375-87. Cuello AC <i>et al.</i> (1980) Use of monoclonal antibodies in immunocytochemistry with special reference to the central nervous system. Brain Res Bull. 5 (5): 575-87. Pioro, E.P. <i>et al.</i> (1985) Loss of substance P immunoreactivity in the nucleus of the spinal trigeminal tract after intradural tumour compression of the trigeminal nerve. Neurosci Lett. 58: 7-12. Mai, J.K. <i>et al.</i> (1986) Substance P in the human brain. Neuroscience. 17 (3): 709-39. Cuello, A.C. <i>et al.</i> (1979) Detection of substance P in the central nervous system by a monoclonal antibody. Proc Natl Acad Sci U S A. 76 (7): 3532-6. Czujkowska, H. & Arciszewski, M.B. (2011) Expression of corticotropin releasing factor (CRF) in the enteric nervous system of the jejunum of sheep Veterinarni medicina 56: 551-60.

7. Kato, A. *et al.* (2012) Endocannabinoid-dependent plasticity at spinal nociceptor synapses. [J Physiol. 590 \(Pt 19\): 4717-33.](#)
8. Arciszewski, M.B. *et al.* (2009) Expression of vasoactive intestinal polypeptide, substance P and neuropeptide Y in jejunal enteric nerves is altered in rabbits suffering from long term *Trichinella spiralis* infection: an immunohistochemical study [Veterinari Medicina, 54: 589–97](#)
9. Bulc, M. *et al.* (2013) Immunohistochemical characterization of the porcine nodose ganglion. [Acta Histochem. 115 \(5\): 440-6.](#)
10. Boorsma, D.M. *et al.* (1982) Direct immunocytochemistry with a horseradish peroxidase-conjugated monoclonal antibody against substance P. [J Histochem Cytochem. 30: 1211-6.](#)
11. Piore, E.P. *et al.* (1984) Loss of substance P and Enkephalin immunoreactivity in the human substantia nigra after striato-pallidal infarction [Brain Res. 292: 339-47.](#)
12. Mortimer, S.T. *et al.* (1990) Immunohistochemical identification of calcitonin gene-related peptide and substance P in nerves of the bovine parathyroid gland. [Cell Tissue Res. 261: 339-45.](#)
13. Galligan, J.J. *et al.* (1988) Changes in surviving nerve fibers associated with submucosal arteries following extrinsic denervation of the small intestine. [Cell Tissue Res. 253: 647-56.](#)
14. Horn, J.P. and Stofer, W.D. (1989) Preganglionic and sensory origins of calcitonin gene-related peptide-like and substance P-like immunoreactivities in bullfrog sympathetic ganglia. [J Neurosci. 9: 2543-61.](#)
15. Verhaert, P. & De Loof, A. (1985) Substance P-like immunoreactivity in the central nervous system of the blattarian insect *Periplaneta americana* L. revealed by a monoclonal antibody. [Histochemistry. 83: 501-7.](#)
16. Polgár, E. *et al.* (2007) A population of large neurons in laminae III and IV of the rat spinal cord that have long dorsal dendrites and lack the neurokinin 1 receptor. [Eur J Neurosci. 26: 1587-98.](#)
17. Nowicki, M. *et al.* (2007) Peptidergic and nitroergic innervation of the pineal gland in the domestic pig: an immunohistochemical study. [Anat Histol Embryol. 36: 311-20.](#)
18. Nair-Roberts, R.G. *et al.* (2006) Distribution of substance P reveals a novel subdivision in the hippocampus of parasitic South American cowbirds. [J Comp Neurol. 496: 610-26.](#)
19. Song, Y. *et al.* (2013) Bilateral increase in expression and concentration of tachykinin in a unilateral rabbit muscle overuse model that leads to myositis. [BMC Musculoskeletal Disord. 14: 134.](#)
20. Szczurkowski, A. *et al.* (2013) Morphology and immunohistochemical characteristics of the pterygopalatine ganglion in the chinchilla (*Chinchilla laniger*, Molina). [Pol J Vet Sci. 16 \(2\): 359-68.](#)
21. Pidsudko, Z. (2014) Immunohistochemical characteristics and distribution of sensory dorsal root Ganglia neurons supplying the urinary bladder in the male pig. [J Mol Neurosci. 52 \(1\): 71-81.](#)
22. Zacharko-Siembida, A. *et al.* (2014) Co-expression patterns of cocaine- and amphetamine-regulated transcript (CART) with neuropeptides in dorsal root ganglia of the pig. [Acta Histochem. 116 \(2\): 390-8.](#)
23. Dudek, A. *et al.* (2012) Immunohistochemical characterization of neurons in the vestibular ganglion (Scarpa's ganglion) of the pig. [Pol J Vet Sci. 15: 499-507.](#)

24. Ibrahim, D. *et al.* (2015) Immunohistochemical studies for the neuronal elements in the vomeronasal organ of the one-humped camel. [J Vet Med Sci. 77 \(2\): 241-5.](#)
25. Christensen, J. *et al.* (2015) Protease-activated receptors in the Achilles tendon-a potential explanation for the excessive pain signalling in tendinopathy. [Mol Pain. 11: 13.](#)
26. Bossowska, A. *et al.* (2015) *Botulinum* Toxin Type A Induces Changes in the Chemical Coding of Substance P-Immunoreactive Dorsal Root Ganglia Sensory Neurons Supplying the Porcine Urinary Bladder. [Toxins \(Basel\). 7 \(11\): 4797-816.](#)
27. Sienkiewicz, W. *et al.* (2015) Innervation of the chinchilla testis, epididymis, and vas deferens [Bull Vet Inst Pulawy. 59 \(4\): 547-56.](#)
28. Czujkowska, A. & Arciszewski, M.B. (2016) Galanin is Co-Expressed with Substance P, Calbindin and Corticotropin-Releasing Factor (CRF) in The Enteric Nervous System of the Wild Boar (*Sus scrofa*) Small Intestine. [Anat Histol Embryol. 45 \(2\): 115-23.](#)
29. Klimczuk, M. *et al.* (2016) Immunohistochemical characterisation of neurons in the mandibular ganglion and nerve fibres supplying the porcine mandibular gland. [Veterinari Medicina. 61: 361-73.](#)
30. Gańko, M. & Całka, J. (2014) Localization and chemical coding of the dorsal motor vagal nucleus (DMX) neurons projecting to the porcine stomach prepyloric area in the physiological state and after stomach partial resection. [J Mol Neurosci. 52 \(1\): 90-100.](#)
31. Pidsudko, Z. *et al.* (2008) Distribution and chemical coding of intramural neurons in the porcine ileum during proliferative enteropathy. [J Comp Pathol. 138 \(1\): 23-31.](#)
32. Kaleczyc, J. *et al.* (2007) The distribution and chemical coding of intramural neurons supplying the porcine stomach - the study on normal pigs and on animals suffering from swine dysentery. [Anat Histol Embryol. 36 \(3\): 186-93.](#)
33. Gańko M & Całka J (2014) Prolonged acetylsalicylic-acid-supplementation-induced gastritis affects the chemical coding of the stomach innervating vagal efferent neurons in the porcine dorsal motor vagal nucleus (DMX). [J Mol Neurosci. 54 \(2\): 188-98.](#)
34. Dudek, A. *et al.* (2017) Chemical Coding of Sensory Neurons Supplying the Hip Joint Capsule in the Sheep. [Anat Histol Embryol. 46 \(2\): 121-31.](#)
35. Zacharko-Siembida, A. *et al.* (2017) An Immunohistochemical Study of Cocaine- and Amphetamine-Regulated Transcript (Cart) Expression in the Pterygopalatine Ganglion of the Pig [Acta Veterinaria. 67 \(3\): 397-408.](#)
36. Rytel, L. *et al.* (2018) Neurochemical characterization of intramural nerve fibres in the porcine oesophagus. [Anat Histol Embryol. 47 \(6\): 517-26.](#)
37. Zalecki, M. (2019) Gastric ulcer induced changes in substance P and Nk1, Nk2, Nk3 receptors expression in different stomach localizations with regard to intrinsic neuronal system. [Histochem Cell Biol. 151 \(1\): 29-42.](#)
38. Zalecki, M. *et al.* (2019) Enteric nervous system in the European beaver (*Castor fiber*) pylorus - an immunohistochemical study. [Pol J Vet Sci. 22 \(1\): 101-7.](#)
39. Sienkiewicz, W. *et al.* (2015) Caudal mesenteric ganglion in the sheep - macroanatomical and immunohistochemical study. [Pol J Vet Sci. 18 \(2\): 379-89.](#)
40. Sickinger, M. *et al.* (2020) Neuropeptides in the urinary tract of male sheep lambs. [Res Vet Sci. 133: 307-12.](#)
41. Szymanska, K. *et al.* (2018) The Influence of High and Low Doses of Bisphenol A (BPA) on the Enteric Nervous System of the Porcine Ileum. [Int J Mol Sci. 19\(3\): 917.](#)
42. Rytel, L. & Całka, J. (2020) Aspirin Administration Affects Neurochemical Characterization of Substance P-Like Immunoreactive (SP-LI) Nodose Ganglia Neurons Supplying the Porcine Stomach. [Biomed Res Int. 2020: 1049179.](#)

43. Palus, K. & Całka, J. (2019) Influence of Acrylamide Administration on the Neurochemical Characteristics of Enteric Nervous System (ENS) Neurons in the Porcine Duodenum. [Int J Mol Sci. 21 \(1\): 15.](#)
44. Makowska, K. & Gonkowski, S. (2021) Changes Caused by Low Doses of Bisphenol A (BPA) in the Neuro-Chemistry of Nerves Located in the Porcine Heart [Animals. 11 \(3\): 780.](#)
45. Bulc, M. *et al.* (2019) Endometritis affects chemical coding of the dorsal root ganglia neurons supplying uterus in the sexually mature gilts. [Res Vet Sci. 124: 417-25.](#)
46. Bulc, M. *et al.* (2018) Changes in Immunoreactivity of Sensory Substances within the Enteric Nervous System of the Porcine Stomach during Experimentally Induced Diabetes. [J Diabetes Res. 2018: 4735659.](#)
47. Palus, K. *et al.* (2019) The impact of low and high doses of acrylamide on the intramural neurons of the porcine ileum. [Food Chem Toxicol. 132: 110673.](#)
48. Meregalli, C. *et al.* (2021) Reversal of Bortezomib-Induced Neurotoxicity by Suvecaltamide, a Selective T-Type Ca-Channel Modulator, in Preclinical Models. [Cancers \(Basel\). 13 \(19\): 5013.](#)
49. Palus, K. *et al.* (2018) Changes in VIP-, SP- and CGRP- like immunoreactivity in intramural neurons within the pig stomach following supplementation with low and high doses of acrylamide. [Neurotoxicology. 69: 47-59.](#)
50. Makowska, K. *et al.* (2022) The Comparison of the Influence of Bisphenol A (BPA) and Its Analogue Bisphenol S (BPS) on the Enteric Nervous System of the Distal Colon in Mice [Nutrients. 15 \(1\): 200.](#)
51. Sienkiewicz, W. *et al.* (2023) Morphology and immunohistochemical characteristics of the otic ganglion in the chinchilla (*Chinchilla laniger* Molina). [Folia Histochem Cytobiol. Jan 24 \[Epub ahead of print\].](#)
52. Benoit, L.B. *et al.* (2023) RBP Image Database: A resource for the systematic characterization of the subcellular distribution properties of human RNA binding proteins. [Nucleic Acids Res. 51 \(D1\): D1549-D1557.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10522 available at: <https://www.bio-rad-antibodies.com/SDS/8450-0505>
10522

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Rat IgG (STAR16...)	DyLight®800
Rabbit Anti Rat IgG (STAR17...)	FITC
Goat Anti Rat IgG (STAR69...)	FITC
Goat Anti Rat IgG (STAR73...)	RPE
Rabbit Anti Rat IgG (STAR21...)	HRP
Goat Anti Rat IgG (MOUSE ADSORBED) (STAR71...)	DyLight®550 , DyLight®650 , DyLight®800
Goat Anti Rat IgG (STAR131...)	Alk. Phos. , Biotin
Goat Anti Rat IgG (STAR72...)	HRP

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M382549:210513'

Printed on 19 Sep 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)