

Datasheet: 8750-2059G BATCH NUMBER 1808

GOAT ANTI CLOSTRIDIUM TETANI TETANUS EXOTOXIN
TETANUS EXOTOXIN
Purified
Polyclonal Antibody
Polyclonal IgG
1 ml

Product Details

Applications	This product has been reported to work in the following applications. This information is						
	derived from testing within our laboratories, peer-reviewed publications or personal						
	communications from the originators. Please refer to references indicated for further						
	information. For general protocol recommendations, please visit <u>www.bio-</u>						
	rad-antibodies.com/protocols.						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry						
	Immunohistology - Frozen						
	Immunohistology - Paraffin			-			
	ELISA	-			1ug/ml		
	Western Blotting	-			1/100 - 1/500		
	Where this product has n	ot been t	ested for u	se in a particular tech	nique this does not		
	necessarily exclude its us	se in such	n procedure	es. Suggested workin	g dilutions are given as		
	a guide only. It is recomn	nended th	at the use	r titrates the product f	or use in their own		
	system using appropriate	negative	/positive c	ontrols.			
		_	-				
Target Species	Bacterial						
Product Form	Purified IgG - liquid						
Antiserum Prenaration	۹ Antisera to C.Tetani Exot	ovin were	raised by	repeated immunication	one of goat with highly		
Antioorani roparation	purified antigen. Purified		-	•			
	purmed antigen. I urmed	igo piepa	area by an	inity chromatography.			
Buffer Solution	Phosphate buffered salin	е					
Preservative	0.09% Sodium Azide (Na	N ₃)					
Stabilisers							
Approx. Protein							
Concentrations	IgG concentration 5.0mg	/ml					

Immunogen	Tetanus Exotoxin.					
External Database Links	UniProt: <u>P04958</u> <u>Related reagents</u>					
RRID	AB_808753					
Specificity	Goat anti tetanus exotoxin antibody recognizes tetanus toxin, also known as tentoxylysin, produced by the soil micro-organism <i>Clostridium tetani</i> . Infection by <i>Clostridium tetani</i> . results in its systemic growth and secretion of tetanus toxin which causes muscle spasms by inhibiting neurotransmitter release. Goat anti tetanus exotoxin antibody also reacts with the C-fragment (~50 kDa carboxy terminal residues 865-1315) of tetanus toxin (<u>Emsley <i>et al.</i> 2000</u>).					
References	1. Paillot, R. <i>et al.</i> (2015) ISCOM-based equine influenza vaccine: Duration of immunity and randomised clinical trials to assess an accelerated schedule of immunisation and efficacy <u>Trials in Vaccinology. 4: 61-70.</u>					
Further Reading	1. Cook, T.M. <i>et al.</i> (2001) Tetanus: a review of the literature. <u>Br J Anaesth. 87 (3):</u> <u>477-87.</u>					
Storage	Store at +4°C or at -20°C if preferred. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.					
Guarantee	12 months from date of despatch					
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/8750-2059G 10040					
Regulatory	For research purposes only					

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) FITC, HRP

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	id.com	Email: antibody_sales_uk@bio-ra	id.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M363528:200528'

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint