

Datasheet: 9100-0004F

Description:	SHEEP ANTI HUMAN TRANSFERRIN:FITC
Specificity:	TRANSFERRIN
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 ml

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u> rad-antibodies.com/protocols.					
		Yes	No	Not Determined	Suggested Dilution	
	Immunofluorescence	•				
	Where this product ha	s not been te	ested for u	se in a particular tec	chnique this does not	
	necessarily exclude its use in such procedures. Suggested working dilutions are given as					
	a guide only. It is reco	mmended the	at the use	r titrates the product	for use in their own	
	system using appropriate negative/positive controls.					
Target Species	Human					
Product Form	Purified IgG - liquid					
Max Ex/Em	Fluorophore	Excitation M	lax (nm)	Emission Max (nm)		
	FITC	490		525		
Antiserum Preparation	i on Antisera to human transferrin were raised by repeated immunisations of sheep with highly purified antigen. Purified IgG prepared by affinity chromatography on protein G					
Buffer Solution	Phosphate buffered saline					
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)					
Approx. Protein Concentrations	IgG concentration 1.0mg/ml					
Immunogen	Pure human transferrin prepared from pooled normal human serum.					

Links P02787 Related reagents P02787 Related reagents Futrez Gene: 7018 TF Related reagents RRID AB_808755 Specificity Sheep anti Human transferrin antibody recognizes human transferrin, an approximately 80 kDa blood plasma glycoprotein synthesised by the liver, which contains two specific high affinity inor (Fe3-b) binding sites and is responsible for the transport an supply of an exchangeable pool of iron, through binding to cell surface transferrin receptors. Transferrin is the primary blood iron transport protein and under normal conditions, approximately one-third of total blood transferrin contains bound iron. Measurement of blood transferrin levels can be used as an indicator for blood iron-carrying capacity and abnormalities of iron metabolism such as anaemia, iron overload and haemochromatosis. Sheep anti Human transferrin antibody shows minimal cross reactivity with related serum proteins. References 1. Olkhov, R.V. & Shaw, A.M. (2014) Growth kinetics of gold nanoparticles on silica/graphene surfaces for multiplex biological immunoassays RSC Adv. 4 (60): 31678-31694. Further Reading 1. Giannetti, A.M. <i>et al.</i> (2003) Mechanism for multiple ligand recognition by the human transferrin receptor. PLoS Biol. 1 (3): E51. S. Rouaut, T.A. (2003) How mammals acquire and distribute iron needed for oxygen-based metabolism. PLoS Biol. 1 (3): E51. S. Rouaut, T.A. (2003) How mammals acquire and distribute iron needed for oxygen-based metabolism. PLoS Biol. 1 (3): E51. S.	External Database				
RND AB_808755 Specificity Sheep anti Human transferrin antibody recognizes human transferrin, an approximately 80 kDa blood plasma glycoprotein synthesised by the liver, which contains two specific high affinity iron (F63+) binding sites and is responsible for the transport and supply of an exchangeable pool of iron, through binding to cell surface transferrin receptors. Transferrin is the primary blood iron transport protein and under normal conditions, approximately one-third of total blood transferrin contains bound iron. Measurement of blood transferrin levels can be used as an indicator for blood iron-carrying capacity and abnormalities of iron metabolism such as anaemia, iron overload and haemochromatosis. Sheep anti Human transferrin antibody shows minimal cross reactivity with related serum proteins. References 1. Okhov, R.V. & Shaw, A.M. (2014) Growth kinetics of gold nanoparticles on silica/graphene surfaces for multiplex biological immunoassays <u>RSC Adv. 4 (60)</u> : 31678-31684. Further Reading 1. Giannetti, A.M. <i>et al.</i> (2003) Mechanism for multiple ligand recognition by the human transferrin receptor. <u>PLoS Biol. 1 (3): E51</u> . 2. Rouault, TA. (2003) How mammals acquire and distribute iron needed for oxygen-based metabolism. <u>PLoS Biol. 1 (3): E79</u> . Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2.8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light. Guarantee 12 months from da	Links	UniProt:			
Tress Related reagents RRID AB_508755 Specificity Sheep anti Human transferrin antibody recognizes human transferrin, an approximately a0 KDa blood plasma glycoprotein synthesised by the liver, which contains two specific synthesized by the liver, which contains the specific synthesized by the liver, which contains the specific synthesized by the liver synthese of gold nanoparticles on sillea/graphene surfaces for multiplex biological immun		P02787 Related reagents			
RRID AB_808755 Specificity Sheep anti Human transferrin antibody recognizes human transferrin, an approximately 80 kDa blood plasma glycoprotein synthesised by the liver, which contains two specific high affinity iron (Fe3+) binding sites and is responsible for the transport and supply of an exchangeable pool of iron, through binding to cell surface transferrin receptors. Transferrin is the primary blood iron transport protein and under normal conditions, approximately one-third of total blood transferrin contains bound iron. Measurement of blood transferrin levels can be used as an indicator for blood foro-carrying capacity and abnormalities of iron metabolism such as anaemia, iron overload and haemochromatosis. Sheep anti Human transferrin antibody shows minimal cross reactivity with related serum proteins. References 1. Olkhov, R.V. & Shaw, A.M. (2014) Growth kinetics of gold nanoparticles on silica/graphene surfaces for multiplex biological immunoassays RSC Adv. 4 (60): 31678-31684. Further Reading 1. Giannetti, A.M. <i>et al.</i> (2003) Mechanism for multiple ligand recognition by the human transferrin receptor. PLoS Biol. 1 (3): E51, 2. Rouault, T.A. (2003) How mammals acquire and distribute iron needed for oxygen-based metabolism. PLoS Biol. 1 (3): E79, Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protecteld from light. <		Entrez Gene:			
Specificity Sheep anti Human transferrin antibody recognizes human transferrin, an approximately 80 kDa blood plasma glycoprotein synthesised by the liver, which contains two specific high affinity iron (Fe3+) binding sites and is responsible for the transport and supply of an exchangeable pool of iron, through binding to cell surface transferrin receptors. Transferrin is the primary blood iron transport protein and under normal conditions, approximately one-third of total blood transferrin contains bound iron. Measurement of blood transferrin levels can be used as an indicator for blood iron-carrying capacity and abnormalities of iron metabolism such as anaemia, iron overload and haemochromatosis. Sheep anti Human transferrin antibody shows minimal cross reactivity with related serum proteins. References 1. Olkhov, R.V. & Shaw, A.M. (2014) Growth kinetics of gold nanoparticles on silica/graphene surfaces for multiplex biological immunoassays RSC Adv. 4 (60): 31678-31684. Further Reading 1. Giannetti, A.M. <i>et al.</i> (2003) Mechanism for multiple ligand recognition by the human transferrin receptor. PLoS Biol. 1 (3): E51. Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light. Guarantee 12 months from date of despatch Health And Safety Materi		7018 TF Related reagents			
80 KDa blood plasma glycoprotein synthesised by the liver, which contains two specific high affinity iron (Fe3+) binding sites and is responsible for the transport and supply of an exchangeable pool of iron, through binding to cell surface transferrin receptors. Transferrin is the primary blood iron transport protein and under normal conditions, approximately one-third of total blood transferrin contains bound iron. Measurement of blood transferrin levels can be used as an indicator for blood iron-carrying capacity and abnormalities of iron metabolism such as anaemia, iron overload and haemochromatosis. Sheep anti Human transferrin antibody shows minimal cross reactivity with related serum proteins. References 1. Olkhov, R.V. & Shaw, A.M. (2014) Growth kinetics of gold nanoparticles on silica/graphene surfaces for multiplex biological immunoassays RSC Adv. 4 (60): 31678-31684. Further Reading 1. Giannetti, A.M. <i>et al.</i> (2003) Mechanism for multiple ligand recognition by the human transferrin receptor. PLoS Biol. 1 (3): E51. 2. Rouault, T.A. (2003) How mammals acquire and distribute iron needed for oxygen-based metabolism. PLoS Biol. 1 (3): E79. Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light. Guarantee 12 months from date of despatch	RRID	AB_808755			
Proteins.References1. Olkhov, R.V. & Shaw, A.M. (2014) Growth kinetics of gold nanoparticles on silica/graphene surfaces for multiplex biological immunoassays RSC Adv. 4 (60): 31678-31684.Further Reading1. Giannetti, A.M. et al. (2003) Mechanism for multiple ligand recognition by the human transferrin receptor. PLoS Biol. 1 (3): E51. 2. Rouault, T.A. (2003) How mammals acquire and distribute iron needed for oxygen-based metabolism. PLoS Biol. 1 (3): E79.StorageThis product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.Guarantee12 months from date of despatchHealth And Safety InformationMaterial Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/9100-0004F 10040	Specificity	80 kDa blood plasma glycoprotein synthesised by the liver, which contains two specific high affinity iron (Fe3+) binding sites and is responsible for the transport and supply of an exchangeable pool of iron, through binding to cell surface transferrin receptors. Transferrin is the primary blood iron transport protein and under normal conditions, approximately one-third of total blood transferrin contains bound iron. Measurement of blood transferrin levels can be used as an indicator for blood iron-carrying capacity and abnormalities of iron metabolism such as anaemia, iron overload and haemochromatosis.			
Silica/graphene surfaces for multiplex biological immunoassays RSC Adv. 4 (60): 31678-31684. Further Reading 1. Giannetti, A.M. et al. (2003) Mechanism for multiple ligand recognition by the human transferrin receptor. PLoS Biol. 1 (3): E51. 2. Rouault, T.A. (2003) How mammals acquire and distribute iron needed for oxygen-based metabolism. PLoS Biol. 1 (3): E79. Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light. Guarantee 12 months from date of despatch Health And Safety Information Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/9100-0004F					
transferrin receptor. PLoS Biol. 1 (3): E51. 2. Rouault, T.A. (2003) How mammals acquire and distribute iron needed for oxygen-based metabolism. PLoS Biol. 1 (3): E79. Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light. Guarantee 12 months from date of despatch Health And Safety Information Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/9100-0004F	References	silica/graphene surfaces for multiplex biological immunoassays RSC Adv. 4 (60):			
-20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light. Guarantee 12 months from date of despatch Health And Safety Information Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/9100-0004F 10040	Further Reading	transferrin receptor. <u>PLoS Biol. 1 (3): E51.</u> 2. Rouault, T.A. (2003) How mammals acquire and distribute iron needed for			
frost-free freezers is not recommended. This product is photosensitive and should be protected from light. Guarantee 12 months from date of despatch Health And Safety Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/9100-0004F	Storage	-20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for			
Health And Safety Material Safety Datasheet documentation #10040 available at: Information https://www.bio-rad-antibodies.com/SDS/9100-0004F 10040		frost-free freezers is not recommended. This product is photosensitive and should be			
Information <u>https://www.bio-rad-antibodies.com/SDS/9100-0004F</u> 10040	Guarantee	12 months from date of despatch			
Regulatory For research purposes only	-	https://www.bio-rad-antibodies.com/SDS/9100-0004F			
	Regulatory	For research purposes only			

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

North & South Tel: +1 800 265 7376 America

Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Europe Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com 'M428025:240301'

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint

Email: antibody_sales_us@bio-rad.com