

Datasheet: AAI21F

BATCH NUMBER 164715

| | |
|----------------------|-----------------------------|
| Description: | SHEEP ANTI BOVINE IgG1:FITC |
| Specificity: | IgG1 |
| Format: | FITC |
| Product Type: | Polyclonal Antibody |
| Isotype: | Polyclonal IgG |
| Quantity: | 1 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | 1/20 - 1/100 |
| Immunohistology - Frozen | ▪ | | | 1/20 - 1/100 |
| Immunohistology - Paraffin | | | ▪ | |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

| Target Species | Bovine | | | | | | |
|-----------------------|--|-------------------|---------------------|-------------------|------|-----|-----|
| Product Form | Purified IgG fraction conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid | | | | | | |
| Max Ex/Em | <table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>525</td> </tr> </tbody> </table> | Fluorophore | Excitation Max (nm) | Emission Max (nm) | FITC | 490 | 525 |
| Fluorophore | Excitation Max (nm) | Emission Max (nm) | | | | | |
| FITC | 490 | 525 | | | | | |

Antiserum Preparation Antisera to bovine IgG1 were raised by repeated immunisation of sheep with highly purified antigen. Purified IgG prepared by affinity chromatography.

Buffer Solution Phosphate buffered saline

Preservative Stabilisers 0.09% Sodium Azide

Approx. Protein Concentrations IgG concentration 1.0 mg/ml

| | |
|--------------------|--|
| Immunogen | Purified bovine IgG1. |
| RRID | AB_323070 |
| Specificity | <p>Sheep anti Bovine IgG1 polyclonal antibody recognizes bovine IgG1.</p> <p>No cross-reactivity with other bovine immunoglobulin classes is seen in immunoelectrophoresis. This product may cross-react with IgG1 from other species.</p> |
| References | <ol style="list-style-type: none"> Vordermeier, H.M. <i>et al.</i> (2003) Improved immunogenicity of DNA vaccination with mycobacterial HSP65 against bovine tuberculosis by protein boosting. Vet Microbiol. 93: 349-59. van Diemen, P.M. <i>et al.</i> (2007) Subunit vaccines based on intimin and Efa-1 polypeptides induce humoral immunity in cattle but do not protect against intestinal colonisation by enterohaemorrhagic <i>Escherichia coli</i> O157:H7 or O26:H-. Vet Immunol Immunopathol. 116: 47-58. von Holtum, C. <i>et al.</i> (2008) Development and evaluation of a recombinant antigen-based ELISA for serodiagnosis of cattle lungworm. Vet Parasitol. 151: 218-26. Patarroyo, J.H. <i>et al.</i> (2009) Immune response of bovines stimulated by synthetic vaccine SBm7462 against <i>Rhipicephalus (Boophilus) microplus</i>. Vet Parasitol. 166: 333-9. Almería, S. <i>et al.</i> (2009) Specific anti-<i>Neospora caninum</i> IgG1 and IgG2 antibody responses during gestation in naturally infected cattle and their relationship with gamma interferon production. Vet Immunol Immunopathol. 130: 35-42. Fiedor, C. <i>et al.</i> (2009) Evaluation of a milk ELISA for the serodiagnosis of <i>Dictyocaulus viviparus</i> in dairy cows. Vet Parasitol. 166: 255-61. Makepeace, B.L. <i>et al.</i> (2009) Immunisation with a multivalent, subunit vaccine reduced patent infection in a natural bovine model of Onchocerciasis during intense field exposure. PLoS Negl. Trop. Dis. 3: e544. Riffault, S. <i>et al.</i> (2010) A new subunit vaccine based on nucleoprotein nanoparticles confers partial clinical and virological protection in calves against bovine respiratory syncytial virus. Vaccine. 28: 3722-34. Ploegaert, T.C. <i>et al.</i> (2010) Genetic variation of natural antibodies in milk of Dutch Holstein-Friesian cows. J Dairy Sci. 93: 5467-73. Van Neerven, R.J. <i>et al.</i> (2010) Milk derived antigen-specific antibodies, methods of preparation and uses thereof. US Patent application no: US20100129377 A1 Colwell, D.D. <i>et al.</i> (2010) <i>Dicrocoelium dendriticum</i> in cattle from Cypress Hills, Canada: Humoral response and preliminary evaluation of an ELISA. Vet Parasitol. 174: 162-5. Prado, M.E. <i>et al.</i> (2011) Vaccination of dairy cows with recombinant <i>Streptococcus uberis</i> adhesion molecule induces antibodies that reduce adherence to and internalization of <i>S. uberis</i> into bovine mammary epithelial cells. Vet Immunol Immunopathol. 141: 201-8. Lavoria, M.Á. <i>et al.</i> (2012) Avidity and subtyping of specific antibodies applied to the indirect assessment of heterologous protection against Foot-and-Mouth Disease Virus in cattle. Vaccine. 30: 6845-50. Assad, A. <i>et al.</i> (2012) Immunophenotyping and characterization of BNP colostrum revealed pathogenic alloantibodies of IgG1 subclass with specificity to platelets, granulocytes and monocytes of all maturation stages. Vet Immunol Immunopathol. 147: 25-34. |

15. Vázquez, L. *et al.* (2012) Antigen-specific antibody isotypes, lymphocyte subsets and cytokine profiles in cattle naturally infested by *Hypoderma sp.* (Diptera: Oestridae). [Vet Parasitol. 184: 230-7.](#)
16. Van Meulder, F. *et al.* (2013) Granule exocytosis of granulysin and granzyme B as a potential key mechanism in vaccine-induced immunity in cattle against the nematode *Ostertagia ostertagi*. [Infect Immun. 81: 1798-809.](#)
17. Mansilla, F.C. *et al.* (2013) Dose-dependent immunogenicity of a soluble *Neospora caninum* tachyzoite-extract vaccine formulated with a soy lecithin/ β -glucan adjuvant in cattle. [Vet Parasitol. pii: S0304-4017\(13\)00252-5.](#)
18. Panadero, R. *et al.* (2013) Effect of reinfestations on systemic immune responses in cattle naturally infested by *Hypoderma sp.* (Diptera: Oestridae). [Vet Parasitol. 193: 238-44.](#)
19. Grit, G.H. *et al.* (2014) Evaluation of cellular and humoral systemic immune response against *Giardia duodenalis* infection in cattle. [Vet Parasitol. 202: 145-55.](#)
20. Maree, F.F. *et al.* (2015) Intra-serotype SAT2 chimeric foot-and-mouth disease vaccine protects cattle against FMDV challenge. [Vaccine. 33 \(25\): 2909-16.](#)
21. Trotta, M. *et al.* (2015) Simultaneous immunization of cattle with foot-and-mouth disease (FMD) and live anthrax vaccines do not interfere with FMD booster responses [Trials in Vaccinology. 4: 38-42.](#)
22. Bautista-Garfias, C.R. *et al.* (2015) Co-immunization of cattle with a vaccine against babesiosis and *Lactobacillus casei* increases specific IgG1 levels to *Babesia bovis* and *B. bigemina*. [Parasitol Int. 64 \(5\): 319-23.](#)
23. González-Hernández A *et al.* (2016) Host protective ASP-based vaccine against the parasitic nematode *Ostertagia ostertagi* triggers NK cell activation and mixed IgG1-IgG2 response. [Sci Rep. 6: 29496.](#)
24. Rybarczyk, J. *et al.* (2017) Effects of lactoferrin treatment on *Escherichia coli* O157:H7 rectal colonization in cattle. [Vet Microbiol. 202: 38-46.](#)
25. Scott, K.A. *et al.* (2017) Evaluation of immune responses of stabilised SAT2 antigens of foot-and-mouth disease in cattle. [Vaccine. 35 \(40\): 5426-33.](#)
26. Rainard, P. *et al.* (2017) Cellular and humoral immune response to recombinant *Escherichia coli*. OmpA in cows. [PLoS One. 12 \(10\): e0187369.](#)
27. Sheng, Z.A. *et al.* (2019) Th2-related cytokines are associated with *Fasciola gigantica* infection and evasion in the natural host, swamp buffalo. [Vet Parasitol. 268: 73-80.](#)
28. Jiménez-Pelayo, L. *et al.* (2019) Early *Neospora caninum* infection dynamics in cattle after inoculation at mid-gestation with high (Nc-Spain7)- or low (Nc-Spain1H)-virulence isolates. [Vet Res. 50 \(1\): 72.](#)
29. Hine, B.C. *et al.* (2019) Immune competence traits assessed during the stress of weaning are heritable and favorably genetically correlated with temperament traits in Angus cattle. [J Anim Sci. 97 \(10\): 4053-4065.](#)
30. Springer, A. *et al.* (2022) Immunization Trials with Recombinant Major Sperm Protein of the Bovine Lungworm *Dictyocaulus viviparus*.. [Pathogens 2022, 11, 55.](#)

Storage

Store at +4°C. DO NOT FREEZE.

This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at:
<https://www.bio-rad-antibodies.com/SDS/AAI21F10040>

Regulatory For research purposes only

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M363602:200528'

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)