

Datasheet: AAI22P

BATCH NUMBER 149395

| Description: | SHEEP ANTI BOVINE IgG2:HRP |
|---------------|----------------------------|
| Specificity: | lgG2 |
| Format: | HRP |
| Product Type: | Polyclonal Antibody |
| Isotype: | Polyclonal IgG |
| Quantity: | 1 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|----------------------|
| Immunohistology - Frozen | | | • | |
| Immunohistology - Paraffin | | | • | |
| ELISA | | | | 1/10,000 - 1/100,000 |
| Western Blotting | | | • | |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

| Target Species | Bovine | |
|-----------------------------------|------------------------------------------------------------------------------------------------------------------------------|-----|
| Product Form | Purified IgG conjugated to Horseradish Peroxidase (HRP) - liq | uid |
| Antiserum Preparation | Antisera to bovine IgG2 were raised by repeated immunisation purified antigen. Purified IgG prepared by affinity chromatogra | |
| Buffer Solution | Phosphate buffered saline | |
| Preservative Stabilisers | 0.05% Proclin™ 300 | |
| Approx. Protein Concentrations | IgG concentration 1.0 mg/ml | |
| Immunogen | Purified bovine IgG2. | |

| RRI | D |
|-----|---|
|-----|---|

AB_323066

Specificity

Sheep anti Bovine IgG2 polyclonal antibody recognizes bovine IgG2.

No cross - reactivity with other bovine immunoglobulin classes is seen in immunoelectrophoresis. This product may cross-react with IgG2 from other species.

References

- 1. Makepeace, B.L. *et al.* (2009) Immunisation with a multivalent, subunit vaccine reduced patent infection in a natural bovine model of Onchocerciasis during intense field exposure. PLoS Negl. Trop. Dis. 3: e544.
- 2. Colwell, D.D. & Goater, C.P. (2010) *Dicrocoelium dendriticum* in cattle from Cypress Hills, Canada: humoral response and preliminary evaluation of an ELISA. <u>Vet Parasitol.</u> 174 (1-2): 162-5.
- 3. Assad, A. *et al.* (2012) Immunophenotyping and characterization of BNP colostra revealed pathogenic alloantibodies of IgG1 subclass with specifity to platelets, granulocytes and monocytes of all maturation stages. <u>Vet Immunol Immunopathol. 147:</u> 25-34.
- 4. Agnes, J.T. *et al.* (2011) Identification of *Anaplasma marginale* Outer Membrane Protein Antigens Conserved between *A. marginale* Sensu Stricto Strains and the Live *A. marginale* subsp. centrale Vaccine Infect Immun. 79: 1311-8.
- 5. Lavoria, M.Á. *et al.* (2012) Avidity and subtyping of specific antibodies applied to the indirect assessment of heterologous protection against Foot-and-Mouth Disease Virus in cattle. <u>Vaccine</u>. 30: 6845-50.
- 6. Vordermeier, H.M. *et al.* (2003) Improved immunogenicity of DNA vaccination with mycobacterial HSP65 against bovine tuberculosis by protein boosting. <u>Vet Microbiol. 93:</u> 349-59.
- 7. Mansilla, F.C. *et al.* (2013) Dose-dependent immunogenicity of a soluble Neospora caninum tachyzoite-extract vaccine formulated with a soy lecithin/β-glucan adjuvant in cattle. <u>Vet Parasitol. 197 (1-2): 13-21.</u>
- 8. Panadero, R. *et al.* (2013) Effect of reinfestations on systemic immune responses in cattle naturally infested by *Hypoderma sp.* (Diptera: Oestridae). <u>Vet Parasitol. 193:</u> 238-44.
- 9. Ploegaert, T.C. *et al.* (2010) Genetic variation of natural antibodies in milk of Dutch Holstein-Friesian cows. J Dairy Sci. 93: 5467-73.
- 10. Van Meulder, F. *et al.* (2013) Granule exocytosis of granulysin and granzyme B as a potential key mechanism in vaccine-induced immunity in cattle against the nematode *Ostertagia ostertagi.* Infect Immun. 81: 1798-809.
- 11. Maree, F.F. *et al.* (2015) Intra-serotype SAT2 chimeric foot-and-mouth disease vaccine protects cattle against FMDV challenge. Vaccine. 33 (25): 2909-16.
- 12. Rybarczyk, J. *et al.* (2017) Effects of lactoferrin treatment on Escherichia coli O157:H7 rectal colonization in cattle. Vet Microbiol. 202: 38-46.
- 13. González-Hernández A *et al.* (2016) Host protective ASP-based vaccine against the parasitic nematode Ostertagia ostertagi triggers NK cell activation and mixed IgG1-IgG2 response. <u>Sci Rep. 6: 29496.</u>
- 14. Pecora, A. *et al.* (2015) Development of an APC-targeted multivalent E2-based vaccine against Bovine Viral Diarrhea Virus types 1 and 2. <u>Vaccine. 33 (39): 5163-71.</u>
- 15. Scott, K.A. et al. (2017) Evaluation of immune responses of stabilised SAT2 antigens

of foot-and-mouth disease in cattle. Vaccine. 35 (40): 5426-33.

16. Sheng, Z.A. *et al.* (2019) Th2-related cytokines are associated with *Fasciola gigantica* infection and evasion in the natural host, swamp buffalo. <u>Vet Parasitol. 268: 73-80.</u>

17. Bucafusco, D .et al. (2019) Immune cells transferred by colostrum do not influence the immune responses to foot-and-mouth disease primary vaccination. <u>J Dairy Sci. 102 (9):</u> 8376-84.

18. Jiménez-Pelayo, L. *et al.* (2019) Early *Neospora caninum* infection dynamics in cattle after inoculation at mid-gestation with high (Nc-Spain7)- or low (Nc-Spain1H)-virulence isolates. Vet Res. 50 (1): 72.

19. Rainard, P. et al. (2017) Cellular and humoral immune response to recombinant *Escherichia coli*. OmpA in cows. PLoS One. 12 (10): e0187369.

20. Springer, A.et al. (2022) Immunization Trials with Recombinant Major Sperm Protein of the Bovine Lungworm *Dictyocaulus viviparus*.. <u>Pathogens 2022, 11, 55.</u>

| Storage | Store at +4°C. DO NOT FREEZE. |
|---------|-------------------------------|
|---------|-------------------------------|

This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.

| Guarantee | 12 months from date of despatch |
|-----------|---------------------------------|
| | |

Acknowledgements Proclin™ 300 is a trademark of The Dow Chemical Company ("Dow") or an affiliated

company of Dow.

Health And Safety Material Safety Datasheet documentation #20392 available at: https://www.bio-rad-antibodies.com/SDS/AAI22P

20392

For research purposes only

Related Products

Regulatory

Recommended Useful Reagents

AbGUARD® HRP STABILIZER PLUS (BUF052A)

AbGUARD® HRP STABILIZER PLUS (BUF052B)

AbGUARD® HRP STABILIZER PLUS (BUF052C)

TMB CORE (BUF056A)

TMB CORE+ (BUF062A)

TMB SIGNAL+ (BUF054A)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M363607:200528'

Printed on 01 Mar 2024