

Datasheet: AAI26F BATCH NUMBER 167357

Description:	GOAT ANTI CAT IgG (Fc):FITC
Specificity:	lgG (Fc)
Format:	FITC
Product Type:	Polyclonal Antibody
lsotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications	derived from testing wi communications from t	thin our laborator he originators. P al protocol recom	in the following application ies, peer-reviewed public ease refer to references i mendations, please visit <u>y</u> Not Determined	ations or personal indicated for further			
	Flow Cytometry		Not Determined	Suggested Dilution			
	Immunohistology - Froze	- -		1/50 - 1/500			
	Immunohistology - Paraff		•	1/30 - 1/300			
	Immunofluorescence		-				
		Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as					
	a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.						
Target Species	Cat						
Species Cross	Reacts with: Iberian Ly	nx, Rabbit					
Reactivity	reactivity is derived fro	m testing within o	nditions may vary betwee ur laboratories, peer-revi nators. Please refer to ref	ewed publications or			
Product Form	Purified IgG conjugate	d to Fluorescein	sothiocyanate Isomer 1 (FITC) - liquid			
Max Ex/Em	Fluorophore	Excitation Max (I	m) Emission Max (nm)				

Antiserum Preparation Antisera to feline IgG were raised by repeated immunisation of goat with highly purified antigen. Purified IgG prepared by affinity chromatography using antigen coupled to agarose beads.

Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide
Stabilisers	0.2% Bovine Serum Albumin
	0.2% Bovine Serum Albumin
Approx. Protein	IgG concentration 1.0 mg/ml
Concentrations	
Immunogen	Purified feline IgG (Fc) fragment.
RRID	AB_323010
Specificity	Goat anti cat IgG (Fc) antibody recognizes feline IgG (Fc) and shows no cross-reactivity with other feline immunoglobulin classes in immunoelectrophoresis.
References	1. Norris, C.R. et al. (2003) Production of polyclonal antisera against feline
	immunoglobulin E and its use in an ELISA in cats with experimentally induced asthma. \underline{Vet}
	Immunol Immunopathol. 96: 149-57.
	2. Jiménez, A. <i>et al.</i> (2008) Membranous glomerulonephritis in the Iberian lynx (<i>Lynx</i>
	pardinus). Vet Immunol Immunopathol. 121: 34-43.
	3. Saari, S. <i>et al.</i> (2008) A nonhealing ulcerative skin lesion associated with <i>Trichinella</i>
	nativa infection in a cat. <u>J Vet Diagn Invest. 20:839-43.</u>
	4. De Craeye, S. <i>et al.</i> (2008) Prevalence of <i>Toxoplasma gondii</i> infection in Belgian house
	cats. <u>Vet Parasitol. 157:128-32.</u>
	5. Tochikura, T.S. <i>et al.</i> (2010) Differential CXCR4 expression and function in
	subpopulations of the feline lymphoma cell line 3201 susceptible to feline
	immunodeficiency virus. <u>J Feline Med Surg. 12: 269-77.</u> 6. Alvarez-Guerrero C <i>et al.</i> (2014) Pathological and parasitological traits in experimentally
	infected cats with <i>Gnathostoma binucleatum</i> (Spirurida: Gnathostomatidae). <u>Vet Parasitol.</u>
	<u>204 (3-4): 279-84.</u>
	7. Persichetti, M.F. <i>et al.</i> (2017) Diagnostic performance of ELISA, IFAT and Western blot
	for the detection of anti- <i>Leishmania infantum</i> antibodies in cats using a Bayesian analysis
	without a gold standard. Parasit Vectors. 10 (1): 119.
	8. Pereira, A. <i>et al.</i> (2019) Antibody response to <i>Phlebotomus perniciosus</i> saliva in cats
	naturally exposed to phlebotomine sand flies is positively associated with Leishmania
	infection. Parasit Vectors. 12 (1): 128.
	9. Cherif, A.M. et al. (2022) A cross- sectional study of Leishmania infantum infection in
	stray cats in Algiers' suburbs, Algeria, and evaluation of serological and molecular tests
	for its diagnosis <u>Veterinaria 71 (1): 73-83.</u>
	10. Balan, P. et al. (2020) Orally administered ovine serum immunoglobulins modulate
	dental plaque in cats. <u>Res Vet Sci. 133: 262-268.</u>
Storage	Store at +4°C. DO NOT FREEZE.
	This product should be stored undiluted. This product is photosensitive and should be
	protected from light. Should this product contain a precipitate we recommend
	microcentrifugation before use.
Guarantos	12 months from data of despetch
Guarantee	12 months from date of despatch

Information	on	https://www.bio 10041	<u>-rad-anti</u>	ibodies.com/SDS/AAI2	<u>6F</u>	
Regulato	ry	For research pu	urposes	only		
	Tel: +1 800 265 73 Fax: +1 919 878 3		ldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50
		ales_us@bio-rad.com		Email: antibody_sales_uk@bio	-rad.com	Email: antibody_sales_de@bio-rad.com

Printed on 12 Apr 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint