Datasheet: AAI29AB

Description:	GOAT ANTI CHICKEN IgG (Fc):Alk. Phos.
Specificity:	IgG (Fc)
Other names:	IgY
Format:	Alk. Phos.
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
ELISA	▪			1/5000 - 1/50000

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

Target Species

Chicken

Product Form

Purified IgG conjugated to Alkaline Phosphatase - liquid

Antiserum Preparation

Antisera to chicken IgG were raised by repeated immunisation of goat with highly purified antigen. Purified IgG prepared by affinity chromatography.

Buffer Solution

50mM HEPES, 0.1M NaCl, 1mM $MgCl_2$, 0.1mM $ZnCl_2$

Preservative Stabilisers

0.09% Sodium Azide
0.2% Bovine Serum Albumin

Approx. Protein Concentrations

IgG concentration 0.5mg/ml

Immunogen

Purified chicken IgG (Fc) fragment.

RRID

AB_10863149

Specificity

Goat anti Chicken IgG (Fc) antibody recognizes the heavy chains of chicken IgG, specifically, epitopes within the Fc region and shows no cross-reactivity with other chicken immunoglobulin classes as assessed by immunoelectrophoresis. This Goat anti Chicken IgG polyclonal antibody does not react with the light chains of chicken IgG.

Goat anti chicken IgG (Fc) has been used successfully for the evaluation of circulating levels of IgG in chickens using ELISA in a number of experimental and field situations.

References

1. Duckworth, J.A. *et al.* (2008) Development of a contraceptive vaccine for the marsupial brushtail possum (*Trichosurus vulpecula*): lack of effects in mice and chickens immunised with recombinant possum ZP3 protein and a possum ZP3 antifertility epitope Wildlife Research 35, 563–72.
2. Peralta, B. *et al.* (2009) Evidence of widespread infection of avian hepatitis E virus (avian HEV) in chickens from Spain. Vet Microbiol. 137: 31-6
3. Norup, L.R. *et al.* (2009) Influence of chicken serum mannose-binding lectin levels on the immune response towards *Escherichia coli.* Poult Sci. 88:543-53.
4. Ferdushy, T. *et al.* (2014) Acquisition of resistance after continuous infection with *Ascaridia galli.* in chickens. Parasitology. : 1-8.
5. Pleidrup, J. *et al.* (2014) *Ascaridia galli* infection influences the development of both humoral and cell-mediated immunity after Newcastle Disease vaccination in chickens. Vaccine. 32 (3): 383-92.
6. Cho Y *et al.* (2015) Proteomic analysis of outer membrane proteins in *Salmonella enterica Enteritidis*. J Microbiol Biotechnol. 25 (2): 288-95.
7. Radomska KA *et al.* (2016) Chicken Immune Response after *In Ovo* Immunization with Chimeric TLR5 Activating Flagellin of *Campylobacter jejuni*. PLoS One. 11 (10): e0164837.
8. Ranchod, H. *et al.* (2018) The antigenicity and cholesteroid nature of mycolic acids determined by recombinant chicken antibodies. PLoS One. 13 (8): e0200298.
9. Vaezirad, M.M. *et al.* (2018) Chicken immune response following in ovo delivery of bacterial flagellin. Vaccine. 36 (16): 2139-46.
10. Naghizadeh, M. *et al.* (2019) Rapid whole blood assay using flow cytometry for measuring phagocytic activity of chicken leukocytes. Vet Immunol Immunopathol. 207: 53-61.
11. Al-karagoly, H. *et al.* (2019) Turkey humoral and cell-mediated immune responses to a Newcastle viscerotropic vaccine and its association with major histocompatibility complex. Bulg J Vet Med. 22 (1): 26-40.
12. Tang, B. *et al.* (2020) GtxA is a virulence factor that promotes a Th2-like response during *Gallibacterium anatis.* infection in laying hens. Vet Res. 51 (1): 40.

Storage

Store at +4°C.
DO NOT FREEZE.

This product should be stored undiluted.
Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

12 months from date of despatch

Health And Safety

Material Safety Datasheet documentation #10089 available at:

Information	https://www.bio-rad-antibodies.com/SDS/AAI29AB 10089
Regulatory	For research purposes only

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	**Worldwide**	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	**Europe**	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M428060:240301'

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint