

Datasheet: AAI30F

BATCH NUMBER 153251

Description:	GOAT ANTI DOG IgM:FITC
Specificity:	IgM
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			1/50 - 1/500
Immunohistology - Paraffin			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

Target Species

Dog

Product Form

Purified IgG fraction conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
FITC	490	525

Antiserum Preparation

Antisera to canine IgM were raised by repeated immunisation of goat with highly purified antigen. Purified IgG prepared by affinity chromatography.

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers

0.09% Sodium Azide
0.2% Bovine Serum Albumin

Approx. Protein Concentrations

IgG concentration 1.0 mg/ml

Immunogen	Purified canine IgM.
External Database Links	UniProt: P01874 Related reagents
RRID	AB_2076336
Specificity	Goat anti Dog IgM polyclonal antibody recognizes canine IgM and shows no cross - reactivity with other canine immunoglobulin classes in immunoelectrophoresis and ELISA. Goat anti Dog IgM may cross react with IgM from other species.
References	<ol style="list-style-type: none"> 1. Peters, I.R. <i>et al.</i> (2004) Measurement of immunoglobulin concentrations in the feces of healthy dogs. Clin Diagn Lab Immunol. 11 (5): 841-8. 2. Carli, E. <i>et al.</i> (2009) Detection of erythrocyte binding IgM and IgG by flow cytometry in sick dogs with Babesia canis canis or Babesia canis vogeli infection. Vet Parasitol. 162: 51-7. 3. Shin, N. <i>et al.</i> (2018) INCB040093 Is a Novel PI3Kδ Inhibitor for the Treatment of B Cell Lymphoid Malignancies. J Pharmacol Exp Ther. 364 (1): 120-30. 4. Sfacteria, A. <i>et al.</i> (2021) Immune Cells and Immunoglobulin Expression in the Mammary Gland Tumors of Dog. Animals (Basel). 11(5):1189. 5. Wyszomtek, M.E. <i>et al.</i> (2022) Canine antibody response against <i>Dirofilaria repens</i>. in natural occult and microfilaremic infections Comparative Immunology, Microbiology and Infectious Diseases. : 101818.
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/AAI30F10041
Regulatory	For research purposes only

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M363635:200528'

Printed on 01 Mar 2024