

Datasheet: AAI31 BATCH NUMBER 160128

Description:	GOAT ANTI DOG IgA
Specificity:	IgA
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .				
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry				
	Immunohistology - Frozen				1/200 - 1/2000
	Immunohistology - Paraffin			•	
	ELISA	-			1/100 - 1/30000
	Immunoprecipitation				
	Western Blotting				Reducing conditions
	Immunodiffusion	-			
	a guide only. It is recomm system using the appropr				for use in their own
Target Species	Dog				
Species Cross	Reacts with: Ferret, Wolf				
Reactivity	Based on sequence similarity, is expected to react with:Mustelid				
	N.B. Antibody reactivity a reactivity is derived from personal communications further information.	testing wi	thin our la	boratories, peer-revie	wed publications or
Product Form	Purified IgG - Liquid				

Antiserum Preparation Antisera to canine IgA were raised by repeated immunisation of goat with highly purified antigen. Purified IgG prepared by affinity chromatography.

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Purified canine IgA.
RRID	AB_323059
Specificity	Goat anti Dog IgA polyclonal antibody recognizes canine IgA and shows no cross reactivity with other canine immunoglobulin classes in immunoelectrophoresis or ELISA.
	Goat anti Dog IgA may cross react with IgA from other species. Goat anti Dog IgA polyclonal antibody has been shown to cross react with IgA from other canids (<u>Frankowiack <i>et al.</i> 2013</u>) and with mustelids (<u>Martel <i>et al.</i> 2009</u>).
References	 Benyacoub, J. <i>et al.</i> (2003) Supplementation of food with <i>Enterococcus faecium</i> (SF68) stimulates immune functions in young dogs. J Nutr. 133 (4): 1158-62. Peters, I.R. <i>et al.</i> (2004) Measurement of immunoglobulin concentrations in the feces of healthy dogs. <u>Clin Diagn Lab Immunol. 11 (5): 841-8.</u> Simpson, K.W. <i>et al.</i> (2009) Influence of <i>Enterococcus faecium</i> SF68 probiotic on giardiasis in dogs. J Vet Intern Med. 23: 476-81. Martel, C.J. & Aasted, B. (2009) Characterization of antibodies against ferret immunoglobulins, cytokines and CD markers. <u>Vet Immunol Immunopathol. 132:109-15.</u> Huang, S.S. <i>et al.</i> (2012) Differential Pathological and Immune Responses in Newly Weaned Ferrets Are Associated with a Mild Clinical Outcome of Pandemic 2009 H1N1 Infection. <u>J Virol. 86: 13187-201.</u> Frankowiack, M. <i>et al.</i> (2013) IgA deficiency in wolves. <u>Dev Comp Immunol. 40 (2): 180-4.</u> Olsson M <i>et al.</i> (2014) The dog as a genetic model for immunoglobulin A (IgA) deficiency: identification of several breeds with low serum IgA concentrations. <u>Vet Immunol Immunopathol. 160 (3-4): 255-9.</u> Frankowiack, M. <i>et al.</i> (2015) IgA deficiency in wolves from Canada and Scandinavia. <u>Dev Comp Immunol. 50 (1): 26-8.</u> Lee, A. <i>et al.</i> (2015) Lack of correlation between mucosal immunoglobulin A-positive plasma cell numbers and TLR5 genotypes in German shepherd dogs reveals association of a locus on CFA 27 with atopic dermatitis. <u>PLoS Genet. 9 (5): e1003475.</u> Martínez-López, L.M. <i>et al.</i> (2021) Hierarchical modelling of immunoglobulin coated bacteria in dogs with chronic enteropathy shows reduction in coating with disease remission but marked inter-individual and treatment-response variability. <u>PLoS One. 16 (8): e0255012.</u> Sfacteria, A. <i>et al.</i> (2021) Immune Cells and Immunoglobulin Expression in the Mammary Gland Tumors of Dog. <u>Animals (Basel). 11(5):1189.</u>

	 13. Rossi, G. <i>et al.</i> (2020) Effects of the Probiotic Mixture Sla Supplement in Healthy Dogs: Evaluation of Fecal Microbiota, Immune Function. <u>Front Vet Sci. 7: 613.</u> 14. Satyaraj, E. <i>et al.</i> (2021) Supplementation of Diets With S and Gut Function in Dogs. <u>Front Nutr. 8: 667072.</u> 				
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Should this product co recommend microcentrifugation before use.	ntain a precipitate we			
Guarantee	12 months from date of despatch				
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/AAI31 10040				
Regulatory	For research purposes only				

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) FITC, HRP

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M363637:200528'

Printed on 10 Jul 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint