

Datasheet: AAI31F

BATCH NUMBER 170522

Description:	GOAT ANTI DOG IgA:FITC
Specificity:	IgA
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Immunohistology - Frozen	▪			1/50 - 1/500
Immunohistology - Paraffin			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

Target Species

Dog

Species Cross Reactivity

Reacts with: Ferret, Wolf

Based on sequence similarity, is expected to react with: Mustelid

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

Purified IgG fraction conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Antiserum Preparation

Antisera to canine IgA were raised by repeated immunisation of goat with highly purified antigen. Purified IgG prepared by affinity chromatography.

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers	0.09% Sodium Azide 0.2% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Purified canine IgA.
Specificity	<p>Goat anti Dog IgA polyclonal antibody recognises canine IgA and shows no cross reactivity with other canine immunoglobulin classes in immunoelectrophoresis or ELISA.</p> <p>Goat anti Dog IgA may cross react with IgA from other species. This anti Dog IgA has been shown to cross react with IgA from other canids (Frankowiack et al. 2013) and with mustelids (Martel et al. 2009).</p>
References	<ol style="list-style-type: none"> 1. Benyacoub, J. <i>et al.</i> (2003) Supplementation of food with <i>Enterococcus faecium</i> (SF68) stimulates immune functions in young dogs. J Nutr. 133 (4): 1158-62. 2. Peters, I.R. <i>et al.</i> (2004) Measurement of immunoglobulin concentrations in the feces of healthy dogs. Clin Diagn Lab Immunol. 11 (5): 841-8. 3. Martel, C.J. & Aasted, B. (2009) Characterization of antibodies against ferret immunoglobulins, cytokines and CD markers. Vet Immunol Immunopathol. 132:109-15. 4. Simpson, K.W. <i>et al.</i> (2009) Influence of <i>Enterococcus faecium</i> SF68 probiotic on giardiasis in dogs. J Vet Intern Med. 23: 476-81. 5. Huang, S.S. <i>et al.</i> (2012) Differential Pathological and Immune Responses in Newly Weaned Ferrets Are Associated with a Mild Clinical Outcome of Pandemic 2009 H1N1 Infection. J Virol. 86: 13187-201. 6. Frankowiack, M. <i>et al.</i> (2013) IgA deficiency in wolves. Dev Comp Immunol. 40 (2): 180-4. 7. Olsson M <i>et al.</i> (2014) The dog as a genetic model for immunoglobulin A (IgA) deficiency: identification of several breeds with low serum IgA concentrations. Vet Immunol Immunopathol. 160 (3-4): 255-9. 8. Frankowiack, M. <i>et al.</i> (2015) IgA deficiency in wolves from Canada and Scandinavia. Dev Comp Immunol. 50 (1): 26-8. 9. Lee, A. <i>et al.</i> (2015) Lack of correlation between mucosal immunoglobulin A-positive plasma cell numbers and TLR5 genotypes in German shepherd dogs with idiopathic chronic enteropathy. J Comp Pathol. 152 (2-3): 201-5. 10. Tengvall, K. <i>et al.</i> (2013) Genome-wide analysis in German shepherd dogs reveals association of a locus on CFA 27 with atopic dermatitis. PLoS Genet. 9 (5): e1003475.
Storage	<p>Store at +4°C. DO NOT FREEZE.</p> <p>This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
Guarantee	12 months from date of despatch.
Health And Safety Information	<p>Material Safety Datasheet documentation available at: https://www.bio-rad-antibodies.com/SDS/AAI31F</p>

Material Safety Datasheet Documentation #10041 available at:

<https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory

For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M256214:140609'

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)