

Datasheet: AAI32

Description:	SHEEP ANTI DOG IgG
Specificity:	IgG
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				
Immunohistology - Frozen	•			1/200 - 1/2000
Immunohistology - Paraffin			•	
ELISA	-			1/100 - 1/30000
Western Blotting	-			1/1000 - 1/30000
Immunodiffusion	-			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

Target Species	Dog	
Species Cross Reactivity	Reacts with: Raccoon dog, Sidestriped Jackal, Black-backed Jackal N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications of personal communications from the originators. Please refer to references indicated for further information.	
Product Form	Purified IgG - liquid	
Antiserum Preparatio	Antisera to canine IgG were raised by repeated immunisation antigen. Purified IgG prepared by affinity chromatography.	of sheep with highly purified
Buffer Solution	Phosphate buffered saline	

Preservative Stabilisers	0.09% Sodium Azide	
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml	
Immunogen	Purified canine IgG.	
RRID	AB_2278376	
Specificity	Sheep anti Dog IgG polyclonal antibody recognizes canine IgG and shows no cross reactivity with other canine immunoglobulin classes in immunoelectrophoresis or ELISA. Sheep anti Dog IgG may cross react with IgG from other species. Cross reactivity has been demonstrated with the Raccoon Dog (<i>Nyctereutes procyonoides</i>), an early canid species (<u>Sukura et al. 2002</u>).	

References

- 1. Benyacoub, J. *et al.* (2003) Supplementation of food with *Enterococcus faecium* (SF68) stimulates immune functions in young dogs. <u>J. Nutr. 133: 1158-1162.</u>
- 2. Xu, L. *et al.* (2007) Immune response after neonatal transfer of a human factor IX-expressing retroviral vector in dogs, cats, and mice. <u>Thromb Res. 120: 269-80.</u>
- 3. Bird, R.C. *et al.* (2011) An autologous dendritic cell canine mammary tumor hybrid-cell fusion vaccine. Cancer Immunol Immunother. 60: 87-97.
- 4. Miller, A.G. (2012) Antiphospholipid antibodies in dogs with immune mediated hemolytic anemia, spontaneous thrombosis, and hyperadrenocorticism. <u>J Vet Intern Med. 26:</u> 614-23.
- 5. Rinkardt, N.E. *et al.* (1999) The effects of prednisone and azathioprine on circulating immunoglobulin levels and lymphocyte subpopulations in normal dogs. <u>Can J Vet Res. 63:</u> 18-24.
- 6. Zhang, J. *et al.* (2004) Neonatal gene transfer with a retroviral vector results in tolerance to human factor IX in mice and dogs. Blood. 103: 143-51.
- 7. Sukura, A. *et al.* (2002) *Trichinella nativa* and *T. spiralis* induce distinguishable histopathologic and humoral responses in the raccoon dog (*Nyctereutes procyonoides*). Vet Pathol. 39: 257-65.
- 8. Carli, E. *et al.* (2009) Detection of erythrocyte binding IgM and IgG by flow cytometry in sick dogs with Babesia canis canis or Babesia canis vogeli infection. <u>Vet Parasitol. 162:</u> 51-7.
- 9. Tan, E. *et al.* (2012) Potentially antigenic RBC membrane proteins in dogs with primary immune-mediated hemolytic anemia. <u>Vet Clin Pathol. 41: 45-55.</u>
- 10. Wieland, B. *et al.* (2012) Prevalence of perinuclear antineutrophilic cytoplasmic autoantibodies in serum of healthy Soft Coated Wheaten Terriers in the United Kingdom. Am J Vet Res. 73: 404-8.
- 11. Xu, L. *et al.* (2003) Neonatal or hepatocyte growth factor-potentiated adult gene therapy with a retroviral vector results in therapeutic levels of canine factor IX for hemophilia B. Blood. 101: 3924-32.
- 12. Spencer, J.A. *et al.* (1999) Presence of antibodies to canine distemper virus, canine parvovirus and canine adenovirus type 1 in free-ranging jackals (Canis adustus and Canis mesomelas) in Zimbabwe. <u>Onderstepoort J Vet Res. 66: 251-3.</u>
- 13. Shapiro, A.J. et al. (2016) Seroprevalence of Coxiella burnetii in Australian dogs.

Zoonoses Public Health. 63 (6): 458-66.

Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	18 months from date of despatch.
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Sheep IgG (H/L) (5184-2304...) Biotin

Donkey Anti Sheep IgG (STAR88...) <u>DyLight®488, HRP</u>

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M339361:181219'

Printed on 09 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint