

## Datasheet: AAI33

<b>Description:</b>	GOAT ANTI HORSE IgM
<b>Specificity:</b>	IgM
<b>Format:</b>	Purified
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	1 mg

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			
Immunoprecipitation			▪	
Western Blotting			▪	
Immunodiffusion	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

<b>Target Species</b>	Horse
<b>Product Form</b>	Purified IgG - liquid
<b>Antiserum Preparation</b>	Antisera to equine IgM were raised by repeated immunisation of goat with highly purified antigen. Purified IgG prepared by affinity chromatography.
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide
<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>Immunogen</b>	Purified equine IgM.
<b>RRID</b>	AB_323035
<b>Specificity</b>	<b>Goat anti horse IgM polyclonal antibody</b> recognizes equine IgM.

No cross-reactivity with other equine immunoglobulin classes is seen in immunoelectrophoresis. This antibody may cross-react with IgM from other species.

---

**References**

1. Jacks, S. *et al.* (2007) Experimental infection of neonatal foals with *Rhodococcus equi* triggers adult-like gamma interferon induction. [Clin Vaccine Immunol. 14: 669-77.](#)
2. Ryan, C. and Giguère, S. (2010) Equine Neonates Have Attenuated Humoral and Cell-Mediated Immune Responses to a Killed Adjuvanted Vaccine Compared to Adult Horses [Clin Vaccine Immunol. 17: 1896-902.](#)
3. Swadzba, M.E. *et al.* (2012) Vitreal IgM Autoantibodies Target Neurofilament Medium in a Spontaneous Model of Autoimmune Uveitis. [Invest Ophthalmol Vis Sci. 53: 294-300.](#)
4. Barrell, E.A. *et al.* (2017) Acute leukemia in six horses (1990-2012). [J Vet Diagn Invest. May 1:1040638717707724 \[Epub ahead of print\]](#)

---

**Storage**

Store at +4°C. DO NOT FREEZE.  
This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee**

12 months from date of despatch

---

**Health And Safety Information**

Material Safety Datasheet documentation #10040 available at:  
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

---

**Regulatory**

For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) [FITC](#), [HRP](#)

**North & South America** Tel: +1 800 265 7376  
Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700  
Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21  
Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M363640:200528'

Printed on 11 Aug 2020