

Datasheet: AA133F

BATCH NUMBER 167622

Description:	GOAT ANTI HORSE IgM:FITC
Specificity:	IgM
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/20 - 1/100
Immunohistology - Frozen	▪			1/100 - 1/1000
Immunohistology - Paraffin			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

Target Species

Horse

Product Form

Purified IgG fraction conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
FITC	490	525

Antiserum Preparation

Antisera to equine IgM were raised by repeated immunisation of goat with highly purified antigen. Purified IgG prepared by affinity chromatography.

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers

0.09% Sodium Azide
0.2% Bovine Serum Albumin

Approx. Protein Concentrations

IgG concentration 1.0 mg/ml

Immunogen	Purified equine IgM.
RRID	AB_323034
Specificity	Goat anti Horse IgM polyclonal antibody recognizes equine IgM. No cross-reactivity with other equine immunoglobulin classes is seen in immunoelectrophoresis. This antibody may cross-react with IgM from other species.
References	<ol style="list-style-type: none"> Jacks, S. <i>et al.</i> (2007) Experimental infection of neonatal foals with <i>Rhodococcus equi</i> triggers adult-like gamma interferon induction. Clin Vaccine Immunol. 14: 669-77. Ryan, C. and Giguère, S. (2010) Equine Neonates Have Attenuated Humoral and Cell-Mediated Immune Responses to a Killed Adjuvanted Vaccine Compared to Adult Horses Clin Vaccine Immunol. 17: 1896-902. Swadzba, M.E. <i>et al.</i> (2012) Vitreal IgM Autoantibodies Target Neurofilament Medium in a Spontaneous Model of Autoimmune Uveitis. Invest Ophthalmol Vis Sci. 53: 294-300. Barrell, E.A. <i>et al.</i> (2017) Acute leukemia in six horses (1990-2012). J Vet Diagn Invest. 29 (4): 529-35. Arroyo, L.G. <i>et al.</i> (2022) Serum IgM antibody response to <i>Clostridioides difficile</i> polysaccharide PS-II vaccination in pony foals. Anaerobe. 77: 102635.
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/AAI33F10041
Regulatory	For research purposes only

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)
'M420518:230706'

Printed on 01 Mar 2024