

Datasheet: AAI38F BATCH NUMBER 148560

Description:	GOAT ANTI HORSE IgG (T):FITC			
Specificity:	lgG (T)			
Format:	FITC			
Product Type:	Polyclonal Antibody			
lsotype:	Polyclonal IgG			
Quantity:	1 mg			

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .							
		Yes	No	Not Determined	Suggested Dilution			
	Flow Cytometry	•			1/20 - 1/100			
	Immunohistology - Frozer	ו ד			1/20 - 1/200			
	Immunohistology - Paraffi	n						
	Where this antibody ha	s not been	tested for u	use in a particular tec	hnique this does not			
	necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.							
Target Species	Horse							
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid							
Max Ex/Em	Fluorophore	Excitation M	lax (nm)	Emission Max (nm)				
	FITC	490		525				
Antiserum Preparatior	i on Antisera to equine IgG (T) were raised by repeated immunisation of goat with highly purified antigen. Purified IgG prepared by affinity chromatography.							
Buffer Solution	Phosphate buffered saline							
Preservative	0.09% Sodium Azide							
Stabilisers	0.2% Bovine Serum Albumin							
Approx. Protein Concentrations	IgG concentration 1.0 r	ng/ml						

Immunogen	Purified equine IgG (T).
RRID	AB_323022
Specificity	Goat anti Horse IgG (T) antibody recognizes equine IgG (T). No cross-reactivity with other equine immunoglobulin classes is seen in immuno-electrophoresis. Goat anti Horse IgG (T) antibody may cross react with IgG from other species.
References	 Hooper-McGrevy, K.E. <i>et al.</i> (2003) Immunoglobulin G subisotype responses of pneumonic and healthy, exposed foals and adult horses to <i>Rhodococcus equi</i> virulence- associated proteins. <u>Clin Diagn Lab Immunol.</u> 10 (3): 345-51. Jacks, S. <i>et al.</i> (2007) Experimental infection of neonatal foals with <i>Rhodococcus equi</i> triggers adult-like gamma interferon induction. <u>Clin Vaccine Immunol.</u> 14: 669-77. Lewis, M.J. <i>et al.</i> (2007) The different effector function capabilities of the seven equine IgG subclasses have implications for vaccine strategies. <u>Mol Immunol.</u> 45: 818-27. Ryan, C. & Giguère, S. (2010) Equine neonates have attenuated humoral and cell-mediated immune responses to a killed adjuvanted vaccine compared to adult horses. <u>Clin Vaccine Immunol.</u> 17 (12): 1896-902. Cauchard S <i>et al.</i> (2014) Assessment of the safety and immunogenicity of <i>Rhodococcus equi</i>-secreted proteins combined with either a liquid nanoparticle (IMS 3012) or a polymeric (PET GEL A) water-based adjuvant in adult horses and foals identification of promising new candidate antigens. <u>Vet Immunol Immunopathol.</u> 157 (3-4): <u>164-74.</u> Meulenbroeks C <i>et al.</i> (2015) Allergen-Specific Cytokine Polarization Protects Shetland Ponies against <i>Culicoides obsoletus</i>-Induced Insect Bite Hypersensitivity. <u>PLoS One.</u> 10 (4): e0122090. Cauchard, S. <i>et al.</i> (2014) Assessment of the safety and immunogenicity of <i>Rhodococcus equi</i>-secreted proteins combined with either a liquid nanoparticle (IMS 3012) or a polymeric (PET GEL A) water-based adjuvant in adult horses and foals identification of promising new candidate antigens. <u>Vet Immunol Immunopathol.</u> 157 (3-4): <u>164-74.</u> Burk, S.V. <i>et al.</i> (2016) Equine antibody response to larval <i>Parascaris equorum</i> excretory-secretory products. <u>Vet Parasitol.</u> 226: 83-7. Lightbody, K.L. <i>et al.</i> (2016) Validation of a novel saliva-based ELISA test for diagnosing tapeworm burden in horses. <u></u>
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/AAI38F 10041

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.com
To find a b	atch/lot specific datash	neet for this produ	uct, please use our online 'M363647:200528'	search tool at	: bio-rad-antibodies.com/datasheets

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint