

# Datasheet: AAI38F

### **BATCH NUMBER 162389**

Description:	GOAT ANTI HORSE IgG (T):FITC
Specificity:	IgG (T)
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

# **Product Details**

### **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/20 - 1/100
Immunohistology - Frozen	•			1/20 - 1/200
Immunohistology - Paraffin				

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

Target Species	Horse			
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liqu		FITC) - liquid	
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)	
	FITC	490	525	

**Antiserum Preparation** Antisera to equine IgG (T) were raised by repeated immunisation of goat with highly purified antigen. Purified IgG prepared by affinity chromatography.

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide 0.2% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml

Immunogen	Purified equine IgG (T).
RRID	AB_323022
Specificity	Goat anti Horse IgG (T) antibody recognizes equine IgG (T). No cross-reactivity with other equine immunoglobulin classes is seen in immuno-electrophoresis.
	Goat anti Horse IgG (T) antibody may cross react with IgG from other species.
References	<ol> <li>Hooper-McGrevy, K.E. et al. (2003) Immunoglobulin G subisotype responses of pneumonic and healthy, exposed foals and adult horses to <i>Rhodococcus equi</i> virulence-associated proteins. Clin Diagn Lab Immunol. 10 (3): 345-51.</li> <li>Jacks, S. et al. (2007) Experimental infection of neonatal foals with <i>Rhodococcus equi</i> triggers adult-like gamma interferon induction. Clin Vaccine Immunol. 14: 669-77.</li> <li>Lewis, M.J. et al. (2007) The different effector function capabilities of the seven equine IgG subclasses have implications for vaccine strategies. Mol Immunol. 45: 818-27.</li> <li>Ryan, C. &amp; Giguère, S. (2010) Equine neonates have attenuated humoral and cell-mediated immune responses to a killed adjuvanted vaccine compared to adult horses. Clin Vaccine Immunol. 17 (12): 1896-902.</li> <li>Cauchard S et al. (2014) Assessment of the safety and immunogenicity of <i>Rhodococcus equi</i>-secreted proteins combined with either a liquid nanoparticle (IMS 3012) or a polymeric (PET GEL A) water-based adjuvant in adult horses and foals-identification of promising new candidate antigens. Vet Immunol Immunopathol. 157 (3-4): 164-74.</li> <li>Meulenbroeks C et al. (2015) Allergen-Specific Cytokine Polarization Protects Shetland Ponies against <i>Culicoides obsoletus</i>-Induced Insect Bite Hypersensitivity. PLoS One. 10 (4): e0122090.</li> <li>Cauchard, S. et al. (2014) Assessment of the safety and immunogenicity of <i>Rhodococcus equi</i>-secreted proteins combined with either a liquid nanoparticle (IMS 3012) or a polymeric (PET GEL A) water-based adjuvant in adult horses and foals-identification of promising new candidate antigens. Vet Immunol Immunopathol. 157 (3-4): 164-74.</li> <li>Burk, S.V. et al. (2016) Equine antibody response to larval <i>Parascaris equorum</i> excretory-secretory products. Vet Parasitol. 226: 83-7.</li> <li>Lightbody, K.L. et al. (2016) Validation of a novel saliva-based ELISA test for diagnosing tapeworm burden in horses. Vet Clin Pathol. 45 (2): 335-46.</li> </ol>
Storage	Store at +4°C. DO NOT FREEZE.  This product should be stored undiluted. This product is photosensitive and should be protected from light.  Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: <a href="https://www.bio-rad-antibodies.com/SDS/AAI38F">https://www.bio-rad-antibodies.com/SDS/AAI38F</a> 10041

# **Regulatory** For research purposes only

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M363647:200528'

### Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint