Datasheet: AAI41P BATCH NUMBER 150791

Description:	GOAT ANTI PIG IgG (Fc):HRP
Specificity:	lgG (Fc)
Format:	HRP
Product Type:	Polyclonal Antibody
lsotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further					
	information. For general protocol recommendations, please visit <u>www.bio-</u>					
	rad-antibodies.com/protocols.					
		Yes	No	Not Determined	Suggested Dilution	
	Immunohistology - Frozen			•		
	Immunohistology - Paraffin			•		
	ELISA				1/10,000 - 1/100,000	
	Western Blotting			•		
	Where this antibody has not been tested for use in a particular technique this does not					
	necessarily exclude its us	se in such	n procedu	res. Suggested workin	g dilutions are given as	
	a guide only. It is recomn	nended th	at the use	er titrates the antibody	for use in their own	
	system using the approp			•		
Target Species	Pig					
Product Form	Purified IgG conjugated t	o Horsera	adish Pero	oxidase (HRP) - liquid		
Antiserum Preparatio	n Antisera to porcine IgG w antigen. Purified IgG pre agarose beads.		• •	-		
Buffer Solution	Phosphate buffered salin	e				
Preservative	0.05% Pro-Clean 400					
Stabilisers	0.2% Bovine Serum Albu	min				
Approx. Protein Concentrations	IgG concentration 1.0 mg	ı/ml				

Immunogen	Purified porcine IgG.
RRID	AB_323036
Specificity	Goat anti Pig IgG (Fc) antibody recognizes the Fc region of the porcine IgG heavy chains and shows no cross - reactivity with other porcine immunoglobulin classes as evaluated by immunoelectrophoresis. Goat anti Pig IgG (Fc) has not been species cross adsorbed and may react with the Fc region of IgG from other species.
	Goat anti Pig IgG (Fc) antibody has been used extensively as a detection reagent for porcine IgG in ELISA, for example monitoring of the IgG response in influenza infected pigs (<u>Crisci <i>et al.</i> 2013</u>).
References	 Assana, E. <i>et al.</i> (2010) Antibody responses to the host-protective <i>Taenia solium</i> oncosphere protein TSOL18 in pigs are directed against conformational epitopes. <u>Parasite Immunol. 32: 399-405.</u> Kick, A.R. <i>et al.</i> (2011) Evaluation of peripheral lymphocytes after weaning and
	vaccination for <i>Mycoplasma hyopneumoniae</i> . <u>Res Vet Sci. 91 (3): e68-72.</u>
	3. Pyo, H. <i>et al.</i> (2010) Serodiagnosis of porcine reproductive and respiratory syndrome virus infection with the use of glycoprotein 5 antigens. <u>Can J Vet Res. 74: 223-7.</u>
	4. Busquets, N. <i>et al.</i> (2010) Experimental infection with H1N1 European swine influenza
	virus protects pigs from an infection with the 2009 pandemic H1N1 human influenza virus.
	<u>Vet Res. 41: 74.</u>
	5. Kang, M.L. et al. (2008) Chitosan microspheres containing Bordetella bronchiseptica
	antigens as novel vaccine against atrophic rhinitis in pigs. <u>J Microbiol Biotechnol. 18:</u>
	<u>1179-85.</u>
	6. Kim, T. <i>et al.</i> (2009) <i>Bordetella bronchiseptica</i> aroA mutant as a live vaccine vehicle for heterologous porcine circovirus type 2 major capsid protein expression. <u>Vet Microbiol.</u>
	<u>138: 318-24.</u>
	7. Picherot, M. <i>et al.</i> (2007) Swine infection with <i>Trichinella spiralis</i> : Comparative analysis
	of the mucosal intestinal and systemic immune responses. Vet Parasitol. 143: 122-30.
	8. Tsai, Y.C. et al. (2010) Porcine circovirus type 2 (PCV2) induces cell proliferation,
	fusion, and chemokine expression in swine monocytic cells <i>in vitro</i> . <u>Vet Res. 41: 60.</u>
	9. Scharek, L. <i>et al.</i> (2005) Influence of a probiotic <i>Enterococcus faecium</i> strain on
	development of the immune system of sows and piglets. <u>Vet Immunol Immunopathol. 105:</u>
	<u>151-61.</u> 10. Scharek, L. <i>et al.</i> (2007) Impact of the probiotic bacteria <i>Enterococcus faecium</i> NCIMB
	10415 (SF68) and <i>Bacillus cereus</i> var. <i>toyoi</i> NCIMB 40112 on the development of serum
	IgG and faecal IgA of sows and their piglets. <u>Arch Anim Nutr. 61: 223-34.</u>
	11. Sheoran, A. et al. (2012) Infection With Cryptosporidium hominis Provides Incomplete
	Protection of the Host Against Cryptosporidium parvum. J Infect Dis 205: 1019-23.
	12. Pina-Pedrero, S. <i>et al.</i> (2012) Genomic and antigenic characterization of monomeric
	autotransporters of <i>Haemophilus parasuis</i> : an ongoing process of reductive evolution. <u>Microbiology. 158: 436-47.</u>
	13. Fleury, A. <i>et al.</i> (2015) <i>Taenia solium</i> : Development of an Experimental Model of
	Porcine Neurocysticercosis. <u>PLoS Negl Trop Dis. 9 (8): e0003980.</u>
	14. Zhang, Y. <i>et al.</i> (2016) Generation of <i>E. coli.</i> -derived virus-like particles of porcine
	circovirus type 2 and their use in an indirect IgG enzyme-linked immunosorbent assay.

	Arch Virol. 161 (6): 1485-91.					
	15. Lee, J.A. et al. (2016) Mycoplasma hyorhinis is a potential pathogen of porcine					
	respiratory disease complex that aggravates pneumonia caused by porcine reproductive					
	and respiratory syndrome virus. Vet Immunol Immunopathol. 177: 48-51.					
	16. Lorenzen E et al. (2017) Intrauterine inoculation of minipigs with Chlamydia					
	trachomatis during diestrus establishes a longer lasting infection compared to vaginal					
	inoculation during estrus. <u>Microbes Infect. Feb 9. pii: S1286-4579(17)30025-4. [Epub</u>					
	ahead of print]					
	17. Zhang, G. et al. (2019) Identification of the B-cell epitopes on N protein of type 2					
	porcine reproductive and respiratory syndrome virus, using monoclonal antibodies. Int J					
	Biol Macromol. 130: 300-6.					
	18. Morales, J. et al. (2018) Persistent Taenia solium. Cysticercosis In the State of					
	Morelos, Mexico: Human and Porcine Seroprevalence. J Parasitol. 104 (5): 465-72.					
	19. Tan, T.K. <i>et al.</i> (2021) A COVID-19 vaccine candidate using SpyCatcher					
	multimerization of the SARS-CoV-2 spike protein receptor-binding domain induces potent					
	neutralising antibody responses. <u>Nat Commun. 12 (1): 542.</u>					
04						
Storage	Store at +4°C. DO NOT FREEZE.					
	This product should be stored undiluted. Should this product contain a precipitate we					
	recommend microcentrifugation before use.					
Guarantee	12 months from date of despatch					
Acknowledgements	Proclin™ 300 is a trademark of The Dow Chemical Company ("Dow") or an affiliated					
	company of Dow.					
Health And Safety	Material Safety Datasheet documentation #20391 available at:					
Information	https://www.bio-rad-antibodies.com/SDS/AAI41P					
	20391					
Regulatory	For research purposes only					

Related Products

Recommended Useful Reagents

AbGUARD® HRP STABILIZER PLUS (BUF052A) AbGUARD® HRP STABILIZER PLUS (BUF052B) AbGUARD® HRP STABILIZER PLUS (BUF052C) TMB CORE (BUF056A) TMB CORE+ (BUF062A) TMB SIGNAL+ (BUF054A)

North & South	Tel: +1 800 265 7376 Worldw	vide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M375009:201211'

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint