

Datasheet: AAI49B

Description:	SHEEP ANTI BOVINE IgA:Biotin
Specificity:	IgA
Format:	Biotin
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen	▪			1/250 - 1/2500
ELISA	▪			1/10,000 - 1/200,000
Immunoprecipitation			▪	
Western Blotting	▪			1/10,000 - 1/200,000
Immunocytochemistry	▪			1/100 - 1/500

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Bovine
Product Form	Purified IgG conjugated to Biotin - liquid
Antiserum Preparation	Antisera to bovine IgA were raised by repeated immunisation of sheep with highly purified antigen. Purified IgG prepared by affinity chromatography.
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide (NaN ₃)
Stabilisers	0.2% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml

Immunogen Purified bovine IgA

Specificity **Sheep anti Bovine IgA antibody** recognizes bovine IgA and shows no cross reactivity with other bovine immunoglobulin classes in immunoelectrophoresis. This antibody may recognize IgA from other species, though this has not been verified.

IgA is the predominant surface secreted immunoglobulin from the intestinal, urogenital and respiratory tracts. IgA functions in a protective capacity, primarily, by preventing viruses and bacteria from adhering to epithelial tissues. High concentrations of IgA are found in saliva, colostrum, milk, tear duct fluid, urine and nasal fluid.

In addition to this polyclonal antibody, we have a monoclonal Mouse anti Bovine IgA , clone IL-A71 antibody ([MCA2438GA](#)) available which is specific for bovine IgA, also available in a peroxidase conjugated format.

References

1. Calvo-Lorenzo, M.S. *et al.* (2016) Wooden hutch space allowance influences male Holstein calf health, performance, daily lying time, and respiratory immunity. [J Dairy Sci. 99 \(6\): 4678-92.](#)
2. González-Hernández A *et al.* (2016) Host protective ASP-based vaccine against the parasitic nematode *Ostertagia ostertagi* triggers NK cell activation and mixed IgG1-IgG2 response. [Sci Rep. 6: 29496.](#)

Further Reading

1. Morton, H.C. *et al.* (2004) Cloning and characterization of an immunoglobulin A Fc receptor from cattle. [Immunology. 111 \(2\): 204-11.](#)

Storage Store at +4°C. DO NOT FREEZE.
This product should be stored undiluted.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at: 10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M339402:181219'

Printed on 22 Sep 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)