

Datasheet: AAM32

| Description: | RABBIT ANTI MOUSE INTERLEUKIN-10 |
|---------------|----------------------------------|
| Specificity: | IL-10 |
| Format: | Purified |
| Product Type: | Polyclonal Antibody |
| Isotype: | Polyclonal IgG |
| Quantity: | 0.1 mg |

Product Details

| Applications | This product has been reported to work in the following applications. This information is | | | | | |
|-----------------------|--|------------|------------|-------------------------|--------------------------|--|
| | derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> . | | | | | |
| | | | | | | |
| | | | | | | |
| | | | | | | |
| | | Yes | No | Not Determined | Suggested Dilution | |
| | Flow Cytometry | | | • | | |
| | Immunohistology - Frozen | | | • | | |
| | Immunohistology - Paraffin | | | • | | |
| | ELISA | - | | | 0.5 - 2.0ug/ml | |
| | Immunoprecipitation | | | • | | |
| | Western Blotting | • | | | 0.1 - 0.2ug/ml | |
| | Functional Assays | • | | | 0.02 - 0.04ug/ml | |
| | Where this antibody has | not been | tested for | use in a particular tec | hnique this does not | |
| Target Species | a guide only. It is recomm system using the appropr Mouse | | | | | |
| Product Form | Purified IgG - lyophilized | | | | | |
| Reconstitution | Reconstitute with 0.1ml distilled water | | | | | |
| | Care should be taken during reconstitution as the protein may appear as a film at the | | | | | |
| | bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution. | | | | | |
| | For long term storage the addition of 0.09% sodium azide is recommended. | | | | | |
| | N.B. For functional studie | | | | | |
| Antiserum Preparation | Antisera to mouse IL-10 | were raise | ed by rep | eated immunisations o | f rabbits with highly | |
| | purified antigen. Purified | lgG was | orepared | from whole serum by a | affinity chromatography. | |
| Buffer Solution | Phosphate buffered salin | ۵ | | | | |

Buffer Solution Phosphate buffered saline

| Preservative Stabilisers | None present |
|-----------------------------------|---|
| Carrier Free | Yes |
| Approx. Protein Concentrations | IgG concentration 1.0mg/ml after reconstitution. |
| Immunogen | Recombinant mouse IL-10. |
| External Database Links | UniProt: <u>P18893</u> <u>Related reagents</u> Entrez Gene: |
| | <u>16153</u> II10 <u>Related reagents</u> |
| Synonyms | II-10 |
| RRID | AB_324032 |
| Specificity | Rabbit anti Mouse Interleukin-10 antibody recognizes mouse Interleukin-10 (IL-10), an immunosuppressive cytokine originally isolated from T helper type 2 cells (Th2), which is also secreted by B lymphocytes, macrophages, monocytes and keratinocytes. IL-10 inhibits the synthesis of several cytokines including Tumour Necrosis Factor, Interferon-gamma and Interleukin-2 and -3, and has a wide range of biological properties, including inhibition of the macrophage and Th cell-mediated cytokine synthesis suppression of the delayed hypersensitivity response. IL-10 may initially be involved in the polarization of Th2 responses. |
| ELISA | AAM32 may be used in an indirect ELISA with <u>PMP21</u> as the standard. |
| Western Blotting | AAM32 may be used under either reducing or non-reducing Western blotting conditions with <u>PMP21</u> as the positive control. |
| References | 1. Ye, Q. <i>et al.</i> (2010) The relationship between collagen scaffold cross-linking agents and neutrophils in the foreign body reaction. <u>Biomaterials. 31 (35): 9192-201.</u> |
| Further Reading | 1. Barrat, F.J. <i>et al.</i> (2002) In vitro generation of interleukin 10-producing regulatory CD4(+) T cells is induced by immunosuppressive drugs and inhibited by T helper type 1 (Th1)- and Th2-inducing cytokines. <u>J Exp Med. 195 (5): 603-16.</u> |
| Storage | Prior to reconstitution store at -20°C. After reconstitution store at -20°C. |
| | This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use. |

| Guarantee | 12 months from date of despatch | | | |
|----------------------------------|--|--|--|--|
| Health And Safety Information | Material Safety Datasheet documentation #10294 available at: https://www.bio-rad-antibodies.com/SDS/AAM32 10294 | | | |
| Regulatory | For research purposes only | | | |

Related Products

Recommended Secondary Antibodies

Sheep Anti Rabbit IgG (STAR34...)FITCGoat Anti Rabbit IgG (H/L) (STAR124...)HRPSheep Anti Rabbit IgG (STAR35...)RPEGoat Anti Rabbit IgG (Fc) (STAR121...)Biotin, FITC, HRPRecommended Positive Controls

RECOMBINANT MOUSE INTERLEUKIN-10 (PMP21)

Recommended Useful Reagents

TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

| North & South | Tel: +1 800 265 7376 | Worldwide | Tel: +44 (0)1865 852 700 | Europe | Tel: +49 (0) 89 8090 95 21 |
|---------------|---------------------------------|-----------|-------------------------------|---------|--------------------------------------|
| America | Fax: +1 919 878 3751 | | Fax: +44 (0)1865 852 739 | | Fax: +49 (0) 89 8090 95 50 |
| | Email: antibody_sales_us@bio-ra | ad.com | Email: antibody_sales_uk@bio- | rad.com | Email: antibody_sales_de@bio-rad.com |

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets M399164:220628'

Printed on 12 Aug 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint