

Datasheet: AAM33 BATCH NUMBER 160289

Description:	GOAT ANTI MOUSE INTERLEUKIN-12
Specificity:	IL-12
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications	This product has been re derived from testing withi communications from the information. For general p rad-antibodies.com/proto	n our lab originato protocol r	oratories, ors. Pleas	peer-reviewed publica e refer to references in	itions or personal ndicated for further
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry			•	
	Immunohistology - Frozen			•	
	Immunohistology - Paraffin			•	
	ELISA	-			0.5 - 2.0 ug/ml
	Immunoprecipitation			•	
	Western Blotting	•			0.1 - 0.2 ug/ml
	Functional Assays				2.0 - 3.0 ng/ml
	necessarily exclude its us a guide only. It is recomm system using the approp	nended th	nat the us	er titrates the antibody	•
Target Species	Mouse				
Product Form	Purified IgG - lyophilized				
Reconstitution	Reconstitute with 0.1ml d Care should be taken du bottom of the vial. Bio-Ra For long term storage the N.B. For functional studie	ring recor ad recomi e addition	nstitution mend that of 0.09%	the vial is gently mixe sodium azide is recon	d after reconstitution.

Antiserum Preparation Antisera to mouse IL-12 were raised by repeated immunisations of goats with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution	Phosphate buffered saline			
Preservative Stabilisers	None present			
Carrier Free	Yes			
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml			
Immunogen	Recombinant mouse IL-12.			
External Database Links	UniProt:P43432Related reagentsP43431Related reagentsEntrez Gene:16160II12bRelated reagents16159II12aRelated reagents			
RRID	AB_323697			
Specificity	Goat anti Mouse Interleukin-12 antibody recognizes mouse interleukin-12 (IL-12). Interleukin-12 is produced by activated B lymphocytes and macrophages. The properties of IL-12 include the stimulation of growth of activated T lymphocytes and the development of TH1 CD4 +ve cells and CD8 +ve cytotoxic T cells. Goat anti Mouse Interleukin-12 antibody neutralizes murine IL-12 activity with an ND ₅₀ for mIL-12 (2ngml ⁻¹ of between 0.002 and 0.002µg ml ⁻¹ .			
ELISA	This purified mouse IL-12 antibody may be used in an indirect ELISA or as the capture reagent in a sandwich ELISA with a <u>biotinylated mouse IL-12 antibody</u> (AAM33B) as the detection reagent.			
Western Blotting	This purified mouse IL-12 antibody may be used in Western Blotting applications under either reducing or non-reducing conditions.			
Storage	Prior to reconstitution store at -20°C. After reconstitution store at -20°C.			
	This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.			
Guarantee	6 months from date of despatch			
Health And Safety	Material Safety Datasheet documentation #10162 available at:			

Informati	<u> </u>	10162	ntibodies.com/SDS/AAM	100	
Regulato	ry F	For research purpose			
Relate	d Products				
Recomm	nended Seco	ndary Antibodies			
Rabbit A	nti Goat IgG (Fe	c) (STAR122) <u>FITC</u>), <u>HRP</u>		
orth & South	Tel: +1 800 265 737	6 Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
orth & South merica	Tel: +1 800 265 737 Fax: +1 919 878 375		Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint