

Datasheet: AAM33B

Description:	GOAT ANTI MOUSE INTERLEUKIN-12:Biotin		
Specificity:	IL-12		
Format:	Biotin		
Product Type:	Polyclonal Antibody		
Isotype:	Polyclonal IgG		
Quantity:	50 µg		

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				
Immunohistology - Frozen				
Immunohistology - Paraffin				
ELISA	-			0.25 - 1.0ug/ml
Western Blotting	-			0.1 - 0.2ug/ml

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

Target Species	Mouse	
Product Form	Purified IgG conjugated to Biotin - lyophilized	
Reconstitution	Reconstitute with 0.5ml sterile PBS containing 0.1% bovine see Care should be taken during reconstitution as the protein may bottom of the vial. Bio-Rad recommend that the vial is gently re-	appear as a film at the
Antiserum Preparation	Antisera to mouse IL-12 were raised by repeated immunisatio purified antigen. Purified IgG was prepared from whole serum	
Buffer Solution	Phosphate buffered saline	
Preservative Stabilisers	None present	

Carrier Free	Yes		
Approx. Protein Concentrations	IgG concentration 0.1mg/ml after reconstitution.		
Immunogen	Recombinant mouse IL-12.		
External Database Links	UniProt: P43432 Related reagents P43431 Related reagents Entrez Gene:		
	16160 II12b Related reagents 16159 II12a Related reagents		
RRID	AB_324598		
Specificity	Goat anti Mouse Interleukin-12 antibody recognizes mouse interleukin-12 (IL-12).		
	Interleukin-12 is produced by activated B lymphocytes and macrophages. The properties of IL-12 include the stimulation of growth of activated T lymphocytes and the development of TH1 CD4 +ve cells and CD8 +ve cytotoxic T cells. Goat anti Mouse Interleukin-12 antibody neutralizes murine IL-12 activity with an ND ₅₀ for		
	mIL-12 (2ngml ⁻¹ of between 0.002 and 0.002μg ml ⁻¹ .		
ELISA	This biotinylated mouse IL-12 antibody may be used in a direct ELISA or as the detection reagent in a sandwich ELISA with a <u>purified mouse IL-12 antibody</u> (AAM33) as the capture reagent.		
Western Blotting	This biotinylated mouse IL-12 antibody may be used in Western Blotting application under either reducing or non-reducing conditions.		
Storage	This product is shipped at ambient temperature. Prior to reconstitution store at -20°C. After reconstitution store at -20°C.		
	This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.		
Guarantee	12 months from date of despatch		
Health And Safety Information	Material Safety Datasheet documentation #10294 available at: https://www.bio-rad-antibodies.com/SDS/AAM33B 10294		
Regulatory	For research purposes only		

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M438979:250523'

Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint