

Datasheet: AAM36

Description:	RABBIT ANTI MOUSE INTERLEUKIN-4
Specificity:	IL-4
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes	No	Not Determined	Suggested Dilution	
	Flow Cytometry					
	Immunohistology - Frozen			•		
	Immunohistology - Paraffin			-		
	ELISA	-			0.5ug/ml	
	Immunoprecipitation			-		
	Western Blotting				0.1ug/ml - 0.2ug/ml	
	Functional Assays				0.25ug/ml - 0.40ug/ml	
	Where this antibody has	not been	tested for	use in a particular te	chnique this does not	
	necessarily exclude its us a guide only. It is recomn system using the approp	nended th	at the use	er titrates the antibody	•	
Target Species	Mouse					
Product Form	Purified IgG - lyophilized					
Reconstitution	Reconstitute with 0.1ml d					
	Care should be taken du	-			•	
	bottom of the vial. Bio-Ra			• •		
	For long term storage the	e addition	of 0.09%	sodium azide is recor	mmendedNB: For	
	functional studies do not	add azide).			
Antiserum Preparatio	n Antisera to mouse IL-4 w		• •		•••	
	purified antigen. Purified	lgG was p	prepared b	by affinity chromatogr	aphy.	
Buffer Solution	Phosphate buffered salin	е				

None present	
Yes	
IgG concentration 1.0 mg/ml after reconstitution.	
Recombinant mouse IL-4	
UniProt:P07750Related reagentsEntrez Gene:16189II4Related reagents	
II-4	
AB_324526	
Rabbit anti Mouse Interleukin-4 antibody recognizes mouse known as B-cell stimulatory factor 1, a 120 amino acid pleiotro 140 amino acid precursor which interacts with multiple cell line fibroblasts and macrophages (<u>Crawford <i>et al.</i> 1987</u>). Interleukin-4 is one of the most important factors involved in B switching (<u>Winter <i>et al.</i> 1987</u>). IL-4 acts as a regulator of low a (CD23) expression on both lymphocytes and monocytes (<u>Miya <i>al.</i> 1987</u>). IL-4 acts through binding to it's cognate receptor CE binds the related cytokine IL-13. Interleukin -4 plays an important role in the alternative activation as indicated by the ability to potently up-regulate the expression macrophage mannose receptor (<u>Stein <i>et al.</i> 1992</u>).	ppic cytokine cleaved from a eages, including T cells, cell IgG1 and IgE class ffinity Fc receptor for IgE ajima <i>et al.</i> 1991, Hudak <i>et</i> 0124, a receptor which also
AAM36 may be used in an indirect ELISA with <u>PMP81</u> as the	standard.
 de Barros, V.E.D. <i>et al.</i> (2011) An experimental model of me Rocio flavivirus in BALB/c mice: inflammatory response, cytok histopathology. <u>Am J Trop Med Hyg. 85 (2): 363-73.</u> Kim, S.R. <i>et al.</i> (2013) Endoplasmic reticulum stress influen pathogenesis by modulating nuclear factor κB activation. <u>J Alle</u> <u>S0091-6749(13)01382-1.</u> 	ine production, and
This product is shipped at ambient temperature. Prior to reconstitution store at -20°C.	
	Yes IgG concentration 1.0 mg/ml after reconstitution. Recombinant mouse IL-4 UniProt: P07750 Related reagents Entrez Gene: 16189 I/4 AB_324526 Rabbit anti Mouse Interleukin-4 antibody recognizes mouse known as B-cell stimulatory factor 1, a 120 amino acid pleiotro 140 amino acid precursor which interacts with multiple cell line fibroblasts and macrophages (Crawford <i>et al.</i> 1987). Interleukin-4 is one of the most important factors involved in B switching (Winter <i>et al.</i> 1987). IL-4 acts as a regulator of low a (CD23) expression on both lymphocytes and monocytes (Miyz <i>al.</i> 1987). IL-4 acts through binding to it's cognate receptor CD binds the related cytokine IL-13. Interleukin -4 plays an important role in the alternative activation as indicated by the ability to potently up-regulate the expression macrophage mannose receptor (Stein <i>et al.</i> 1992). AAM36 may be used in an indirect ELISA with PMP81 as the 1. de Barros, V.E.D. <i>et al.</i> (2011) An experimental model of mor Rocio flavivirus in BALB/c mice: inflammatory response, cytok histopathology. Am J Trop Med Hyg. 85 (2): 363-73. 2. Kim, S.R. <i>et al.</i> (2013) Endoplasmic reticulum stress influer

	This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.		
Guarantee	12 months from date of despatch		
Health And Safety Information	Material Safety Datasheet documentation #10294 available at: https://www.bio-rad-antibodies.com/SDS/AAM36 10294		
Regulatory	For research purposes only		

Related Products

Recommended Secondary Antibodies

Goat Anti Rabbit IgG (H/L) (STAR124...) <u>HRP</u>

Sheep Anti Rabbit IgG (STAR35...) RPE

Goat Anti Rabbit IgG (Fc) (STAR121...) Biotin, FITC, HRP

Recommended Useful Reagents

TidyBlot WESTERN BLOT DETECTION REAGENT:HRP (STAR209P)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	d.com	Email: antibody_sales_uk@bio-ra	ad.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M438983:250523'

Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint