

## Datasheet: AAM57B

RABBIT ANTI MOUSE SF-20:Biotin
SF-20
INTERLEUKIN-25
Biotin
Polyclonal Antibody
Polyclonal IgG
50 μg

## **Product Details**

## **Applications**

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <a href="www.bio-rad-antibodies.com/protocols">www.bio-rad-antibodies.com/protocols</a>.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry				
Immunohistology - Frozen				
Immunohistology - Paraffin				
ELISA	•			0.25 - 1.0ug/ml
Immunoprecipitation			•	
Western Blotting	•			0.1 - 0.2ug/ml

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Mouse	
Product Form	Purified IgG conjugated to Biotin - lyophilized	
Reconstitution	Reconstitute with 0.5ml sterile PBS containing 0.1% Bovine Sociate should be taken during reconstitution as the protein may bottom of the vial. Bio-Rad recommend that the vial is gently nNB: For long term storage the addition of 0.09% sodium azide	appear as a film at the nixed after reconstitution.
Antiserum Preparation	Antisera to mouse SF-20 were raised by repeated immunisation purified antigen. Purified IgG was prepared by affinity chromat	<b>,</b>
Buffer Solution	Phosphate buffered saline	

Preservative Stabilisers	None present
Carrier Free	Yes
Immunogen	Recombinant mouse SF-20.
External Database Links	UniProt:  Q9CPT4 Related reagents  Entrez Gene:
	28106 D17Wsu104e Related reagents
Synonyms	II25
RRID	AB_2124934
Specificity	<b>Rabbit anti Mouse SF-20 antibody</b> recognizes mouse stromal cell-derived growth factor 20 (SF-20), otherwise known as interleukin-25 (IL-25), a 166 amino acid secreted growth factor and member of the UPF0556 family.
	SF-20 should not be confused with interleukin 17E (IL-17E), which is also alternatively called IL-25.
Storage	Prior to reconstitution store at -20°C.  After reconstitution store at -20°C.
	This product should be stored undiluted. Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10294 available at: <a href="https://www.bio-rad-antibodies.com/SDS/AAM57B">https://www.bio-rad-antibodies.com/SDS/AAM57B</a> 10294
Regulatory	For research purposes only

America

North & South Tel: +1 800 265 7376

Worldwide

Tel: +44 (0)1865 852 700

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +1 919 878 3751

Email: antibody\_sales\_us@bio-rad.com

Fax: +44 (0)1865 852 739

Fax: +49 (0) 89 8090 95 50

Email: antibody\_sales\_uk@bio-rad.com

Email: antibody\_sales\_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M402131:220718'

Printed on 12 Aug 2023